

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
29 June 2006 (29.06.2006)

PCT

(10) International Publication Number  
**WO 2006/069153 A2**

(51) International Patent Classification:  
*C07C 65/40* (2006.01) *A61K 31/19* (2006.01)  
*C07C 65/24* (2006.01) *A61P 3/14* (2006.01)  
*A61K 31/165* (2006.01)

(21) International Application Number:  
PCT/US2005/046360

(22) International Filing Date:  
19 December 2005 (19.12.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/637,930 21 December 2004 (21.12.2004) US

(71) Applicant (for all designated States except US): **ELI LILLY AND COMPANY** [US/US]; Lilly Corporate Center, Indianapolis, Indiana 46285 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GOSSETT, Lynn, Stacy** [US/US], 7345 Donegal Drive, Indianapolis, Indiana 46217 (US). **LOPEZ, Jose, Eduardo** [DO/US], 10224 Cheswick Lane, Fishers, Indiana 46038 (US). **WARSHAWSKY, Alan, M.** [US/US], 13828 Springmill Ponds Circle, Carmel, Indiana 46032 (US). **YEE, Ying, Kwong** [US/US], 5127 Briarstone Trace, Carmel, Indiana 46033 (US).

(74) Agents: **MYERS, James, B.** et al.: P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**


- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

**Published:**

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: VITAMIN D RECEPTOR MODULATORS



(57) Abstract: The present invention relates to novel, non-secosteroidal, phenyl-naphthalene compounds of Formula (I) wherein R, R<sub>1</sub>, RP, Z<sub>p</sub>, L<sub>p1</sub>, L<sub>p2</sub>, L<sub>NP</sub>, RP<sub>3</sub>, RN, and Z<sub>NP</sub> are defined herein, their preparation, pharmaceutical compositions, and methods of use.

-1-

## VITAMIN D RECEPTOR MODULATORS

### REFERENCE TO RELATED APPLICATION

5           This application claims the benefit of priority under title 35 United States Code § 119(e) of Provisional Patent Application No. 60/637,930 filed December 21, 2004, which is incorporated herein by reference in its entirety

### BACKGROUND OF THE INVENTION

10           Vitamin D Receptor (VDR) is a ligand dependent transcription factor that belongs to the superfamily of nuclear hormone receptors. The VDR protein is 427 amino acids, with a molecular weight of ~50 kDa. The VDR ligand,  $1\alpha,25$ -dihydroxyvitamin D3 (the hormonally active form of Vitamin D) has its action mediated by its interaction with the  
15           nuclear receptor known as Vitamin D receptor ("VDR"). The VDR ligand,  $1\alpha,25$ -dihydroxyvitamin D3 ( $1\alpha,25(\text{OH})_2\text{D}_3$ ) acts upon a wide variety of tissues and cells both related to and unrelated to calcium and phosphate homeostasis.

          The activity of  $1\alpha,25$ -dihydroxyvitamin D3 in various systems suggests wide clinical applications. However, use of conventional VDR ligands is hampered by their  
20           associated toxicity, namely hypercalcemia (elevated serum calcium). Currently,  $1\alpha,25(\text{OH})_2\text{D}_3$ , marketed as Rocaltrol® pharmaceutical agent ( product of Hoffmann-La Roche), is administered to kidney failure patients undergoing chronic kidney dialysis to treat hypocalcemia and the resultant metabolic bone disease. Other therapeutic agents, such as Calcipotriol® (synthetic analog of  $1\alpha,25(\text{OH})_2\text{D}_3$  ) show increased separation of  
25           binding affinity on VDR from hypercalcemic activity.

          Chemical modifications of  $1\alpha,25(\text{OH})_2\text{D}_3$  have yielded analogs with attenuated calcium mobilization effects (R. Bouillon et. al., Endocrine Rev. 1995, 16, 200-257). One such analog, Dovonex ® pharmaceutical agent (product of Bristol-Meyers Squibb Co.), is currently used in Europe and the United States as a topical treatment for mild to moderate  
30           psoriasis (K. Kragballe et. al., Br. J. Dermatol. 1988, 119, 223-230).

-2-

Other Vitamin D<sub>3</sub> mimics have been described in the publication, Vitamin D Analogs: Mechanism of Action of Therapeutic Applications, by Nagpal, S.; Lu, J.; Boehm, M. F., Curr. Med. Chem. 2001, 8, 1661-1679.

Although some degree of separation between the beneficial action and calcium  
5 raising (calcemic) effects has been achieved with these VDR ligands, to date the separation has been insufficient to allow for oral administration to treat conditions such as osteoporosis, cancers, leukemias, and severe psoriasis.

One example of a major class of disorder that could benefit from VDR mediated biological efficacy in the absence of hypercalcemia is osteoporosis. Osteoporosis is a  
10 systemic disorder characterized by decreased bone mass and microarchitectural deterioration of bone tissue leading to bone fragility and increased susceptibility to fractures of the hip, spine, and wrist (World Health Organization WHO 1994). Osteoporosis affects an estimated 75 million people in the United States, Europe, and Japan.

15 Within the past few years, several antiresorptive therapies have been introduced. These include bisphosphonates, hormone replacement therapy (HRT), a selective estrogen receptor modulator (SERM), and calcitonins. These treatments reduce bone resorption, bone formation, and increase bone density. However, none of these treatments increase true bone volume nor can they restore lost bone architecture.

20 Another major disorder that could benefit from VDR mediated biological activity is psoriasis. Psoriasis is one of the most common dermatologic diseases and is a chronic inflammatory skin condition characterized by erythematous, sharply demarcated papules and rounded plaques, covered by silvery micaceous scale.

Synthetic VDR ligands with reduced calcemic potential have been synthesized.  
25 For example, a class of bis-phenyl compounds stated to mimic 1 $\alpha$ , 25-dihydroxyvitamin D<sub>3</sub> is described in US Patent No. 6,218,430 and the article; "Novel nonsecosteroidal vitamin D mimics exert VDR-modulating activities with less calcium mobilization than 1 $\alpha$ , 25-Dihydroxyvitamin D<sub>3</sub>" by Marcus F. Boehm, et. al., Chemistry & Biology 1999, Vol 6, No. 5, pgs. 265-275.

30 Synthetic VDR ligands having an aryl-thiophene nucleus are described in United States provisional patent application SN 60/384,151, filed 29 May 2002 (WO


-3-

03/101,978), and synthetic VDR ligands having phenyl-benzoxazole nucleus are described in United States provisional patent application SN 60/638,029 filed 21 December 2004.


There remains a need for improved treatments using alternative or improved pharmaceutical agents that mimic  $1\alpha, 25$ -dihydroxyvitamin  $D_3$  to stimulate bone formation, restore bone quality, and treat other diseases without the attendant disadvantage of hypercalcemia.

### SUMMARY OF THE INVENTION

Novel compounds having a phenyl-naphthalene nucleus of Formula "(PN)" have been found effective as Vitamin D Receptor modulators (VDRM):



Compounds of the invention with VDR modulating activities are represented by Formula I



wherein the variables R, R', RP, RP<sub>3</sub>, L<sub>P1</sub>, L<sub>P2</sub>, Z<sub>P</sub>, RN, L<sub>NP</sub>, and Z<sub>NP</sub> are as hereinafter defined.

In another aspect, the present invention is directed towards pharmaceutical compositions containing pharmaceutically effective amounts of compounds of Formula I

-4-

or a pharmaceutically acceptable salt or a prodrug thereof, either singly or in combination, together with pharmaceutically acceptable carriers and/or auxiliary agents.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of osteoporosis containing pharmaceutically effective amounts of the vitamin D receptor modulator compound of Formula I alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of osteoporosis.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of psoriasis containing pharmaceutically effective amounts of the vitamin D receptor modulator compound of Formula I alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of psoriasis.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of prostate cancer containing pharmaceutically effective amounts of the vitamin D receptor modulator compound of Formula I alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of prostate cancer.

Another aspect of the invention is to use the compounds of Formula I to treat disease states responsive to Vitamin D receptor ligands.

Another aspect of the invention is the prevention and treatment of acne, actinic keratosis, alopecia, Alzheimer's disease, autoimmune induced diabetes, benign prostatic hyperplasia, bladder cancer, bone fracture healing, breast cancer, Crohn's disease, prostate cancer, colon cancer, Type I diabetes, host-graft rejection, hypercalcemia, Type II diabetes, leukemia, multiple sclerosis, insufficient sebum secretion, osteomalacia, osteoporosis, insufficient dermal firmness, insufficient dermal hydration, myelodysplastic syndrome, psoriatic arthritis, psoriasis, renal osteodystrophy, rheumatoid arthritis, scleroderma, seborrheic dermatitis, skin cancer, systemic lupus erythematosus, skin cell damage from mustard vesicants, ulcerative colitis, and wrinkles, by administering to a mammal in need thereof a pharmaceutically effective amount of a compound of Formula I.

## DETAILED DESCRIPTION OF THE INVENTION

Definitions:

-5-

The term, "abscess" refers to adverse complications often associated with surgery, trauma, or diseases that predispose the host to abscess formation from encapsulated bacteria lymphocytes, macrophages, and etc.


The term, "adhesion" refers to the adverse and abnormal union of surfaces normally separate by the formation of new fibrous tissue resulting from an inflammatory process.

The term, "compound of the invention" refers to a compound represented by Formula I or as set out as products of the Examples or synthesis schemes described herein.


The term, "Active Ingredient" means a compound of the invention.

The term, "mustard" is inclusive of both sulfur mustards and nitrogen mustards, either alone or in any combination. Exemplary of such compounds are the vesicants; bis(2-chloroethyl) sulfide (Chemical Agent Symbol HD),  $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$  1,2-bis(2-chloroethylthio)ethane (Chemical Agent Symbol Q),  $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$ ; bis(2-chloroethylthioethyl) ether,  $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)\text{O}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$  (Chemical Agent Symbol T); tris(2-chloroethyl) amine (Chemical Agent Symbol HN3)  $\text{N}(\text{CH}_2\text{CH}_2\text{Cl})_3$ ; N-methyl-2,2'-dichlorodiethylamine (Chemical Agent Symbol NH2); and 2,2'-dichlorotriethylamine,  $\text{CH}_3\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2\text{Cl})_2$  (Chemical Agent Symbol NH1).

The term heteroaryl as used herein refers to the heteroaryls illustrated below:



-6-



-7-

where the dotted line crossing a solid line symbol represents a bond of attachment between the atom of the radical and the rest of the molecule.

The term, "(Acidic Group)" means an organic group that acts as a proton donor capable of hydrogen bonding. Illustrative of an (Acidic Group) is a group selected from the following: carboxylic acid, acylsulfonamide, tetrazolyl, substituted heteroaryls with acidic hydrogens, i.e., hydroxyl groups.

The term, "mammal" includes humans.

The terms "halo" and halogen refer to fluorine, chlorine, bromine, and iodine. Preferred halogens for the present invention include fluorine.

Unless specified herein, chemical terms are used in their customary usage as understood by one skilled in the art.

The term, "C<sub>1-3</sub> alkyl" refers to an alkyl group selected from methyl, ethyl, n-propyl, and isopropyl. The abbreviations, "Me" means methyl; "Et" means ethyl; "iPr" or "i-Pr" means 1-methylethyl; and "tBu" or "t-Bu" means 1,1-dimethylethyl.

The term, "branched C<sub>3</sub>-C<sub>5</sub> alkyl" is an alkyl group selected from 1-methylethyl; 1-methylpropyl; 2-methylpropyl; 1,1-dimethylethyl; 1,1-dimethylpropyl; 1,2-dimethylpropyl; or 2,2-dimethylpropyl. Preferred branched C<sub>3</sub>-C<sub>5</sub> alkyl groups are 2-methylpropyl and 1,1-dimethylethyl, with the 1,1-dimethylethyl group being most preferred.

The term "alkenyl" refers to aliphatic groups wherein the point of attachment is a carbon-carbon double bond, for example vinyl, 1-propenyl, and 1-cyclohexenyl. Alkenyl groups may be straight-chain, branched-chain, cyclic, or combinations thereof, and may be optionally substituted. It will be understood that alkenyl groups can include one or more double bonds. Further, the alkenyl groups can include positional isomers about the double bonds i.e. trans (Z) or cis (E) isomers. Suitable alkenyl groups have from 2 to about 20 carbon atoms. It also will be understood by those skilled in the art that compounds of the present invention can exist in two or more tautomeric forms. All such tautomeric forms are contemplated to be included within the scope of the present invention.

The term "C<sub>1</sub>-C<sub>5</sub> alkyl" refers to saturated aliphatic groups including straight-chain, branched-chain, and cyclic groups and any combinations thereof. Examples of C<sub>1</sub>-



-8-

C<sub>5</sub> alkyl groups are methyl, ethyl, n-propyl, from 1-methylethyl; n-butyl, 1-methylpropyl; 2-methylpropyl; 1,1-dimethylethyl; n-amyl, 1,1-dimethylpropyl; 1,2-dimethylpropyl; and 2,2-dimethylpropyl.


The term "cycloalkyl" includes organic radicals having 3 to 8 carbon atoms as  
 5 ring members. Examples include: cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. When substituted, the substituents can be selected from halo, hydroxyl, -CN, C<sub>1</sub>-C<sub>3</sub> alkyl, -SH, -OC<sub>1</sub>-C<sub>3</sub> alkyl, and -SC<sub>1</sub>-C<sub>3</sub> alkyl.

The term, "cycloalkenyl" includes organic radicals having 3 to 8 carbon atoms as  
 ring members; non-limiting examples include: cyclopropenyl, cyclobutenyl,  
 10 cyclopentenyl and cyclohexenyl.

The term, "C<sub>1</sub>-C<sub>5</sub> haloalkyl" is an alkyl group containing one or more halogen  
 atoms. The term, "C<sub>1</sub>-C<sub>5</sub> fluoroalkyl" is an alkyl group containing fluorine and includes  
 organic radicals such as -CF<sub>3</sub>, -CHF<sub>2</sub>, -CH<sub>2</sub>F, -CF<sub>2</sub>CF<sub>3</sub>, -CHF<sub>2</sub>CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub>,  
 -CH<sub>2</sub>CHF<sub>2</sub>, and -CH<sub>2</sub>CH<sub>2</sub>F, with -CF<sub>3</sub> being preferred.

The term, "hydroxyalkyl" means an alkyl group having at least one hydroxyl  
 group. Non-limiting examples include: 3-methyl-3-hydroxypentyl, 3-methyl-3-  
 hydroxypentenyl, 3-methyl-3-hydroxypentynyl, 3-ethyl-3-hydroxypentyl, 3-ethyl-3-  
 hydroxypentenyl, 3-ethyl-3-hydroxypentynyl, 3-ethyl-3-hydroxy-4-methylpentyl, 3-ethyl-  
 3-hydroxy-4-methylpentenyl, 3-ethyl-3-hydroxy-4-methylpentynyl, 3-propyl-3-  
 20 hydroxypentyl, 3-propyl-3-hydroxypentenyl, 3-propyl-3-hydroxypentynyl, 1-hydroxy-2-  
 methyl-1-(methylethyl)propyl, 2-methyl-3-hydroxy-4,4-dimethylpentyl, 2-methyl-3-  
 hydroxy-3-ethylpentyl, 2-ethyl-3-hydroxy-3-ethylpentyl, 2-ethyl-3-hydroxy-4,4-  
 dimethylpentyl, 1-hydroxycycloalkenyl; and 1-hydroxycycloalkyl.

The term "hydroxycycloalkyl" refers to a radical having the general structural  
 25 formula:




where w is an integer from 1 to 6 and the hydroxyl radical is substituted on any ring  
 carbon atom. Examples include: 2-hydroxycyclohexylmethyl, 3-methyl-2-

-9-


hydroxycyclohexyloxy, 3-methyl-2-hydroxycyclohexylmethyl, and 3,3-dimethyl-2-hydroxycyclohexyloxy.

The term "1-hydroxycycloalkyl" refers to a radical having the general structural formula:



where  $w$  is defined as above. Examples of 1-hydroxycycloalkyl radicals include: 1-hydroxycyclopropyl, 1-hydroxycyclobutyl, 1-hydroxycyclopentyl, 1-hydroxycyclohexyl, 1-hydroxycycloheptyl, and 1-hydroxycyclooctyl.

The term oxocycloalkyl refers to a radical having the general structural formula:



where  $w$  is defined as above. The bond of attachment of the oxocycloalkyl to the referenced molecule need not be restricted to the carbon adjacent to the carbonyl carbon, but can be attached via any of the carbon atoms making up the rings. Non-limiting examples of oxocycloalkyl radicals include: 2-oxocyclohexyloxy, 2-oxocyclohexylmethyl, 3-methyl-2-oxocyclohexyloxy, 3-methyl-2-oxocyclohexylmethyl, 3,3-dimethyl-2-oxocyclohexyloxy, 3,3-dimethyl-2-oxocyclohexylmethyl, and 2-hydroxycyclohexyloxy.

Certain compounds of the invention exist in isomeric configurations with chiral centers, i.e., diastereomers and enantiomers. Each of the isomeric forms of the compounds is contemplated to be within the scope of the present invention. Each of the various isomers can be prepared as single isomers and/or separated into single isomers by techniques known to those skilled in the art. Therefore, the compounds of the present invention can be used either as single isomer or isomeric form or alternatively the compounds of the present invention can be used as a combination of isomers. The

-10-


“jagged” bond illustrated below is used to represent that carbon to which it is attached can exist as either configuration, i.e., R or S.



- 5 It also will be understood by those skilled in the art that compounds of the present invention can exist in two or more tautomeric forms. All such tautomeric forms are contemplated to be included within the scope of the present invention.

#### Compounds of the Invention:

- 10 The compounds of the invention with vitamin receptor modulating (VDRM) activity are represented by Formula (I) or a pharmaceutically acceptable salt or a prodrug derivative thereof:



- 15 wherein

R and R' are independently C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, or together R and R' form a substituted or unsubstituted, saturated or unsaturated cycloalkyl ring having from 3 to 8 carbon atoms;

- 20 RP<sub>3</sub> and RN are independently selected from the group consisting of hydrogen, halo, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, -O-C<sub>1</sub>-C<sub>5</sub> alkyl, -S-C<sub>1</sub>-C<sub>5</sub> alkyl, -O-C<sub>1</sub>-C<sub>5</sub> haloalkyl, -CN, -NO<sub>2</sub>, acetyl, -S-C<sub>1</sub>-C<sub>5</sub> haloalkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, and C<sub>3</sub>-C<sub>5</sub> cycloalkenyl;

-11-

RP is selected from: hydrogen, halo, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, -O-C<sub>1</sub>-C<sub>5</sub> alkyl, -S-C<sub>1</sub>-C<sub>5</sub> alkyl, -O-C<sub>1</sub>-C<sub>5</sub> haloalkyl, -CN, -NO<sub>2</sub>, acetyl, -S-C<sub>1</sub>-C<sub>5</sub> haloalkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, and C<sub>3</sub>-C<sub>5</sub> cycloalkenyl;

(L<sub>P1</sub>), (L<sub>P2</sub>), and (L<sub>NP</sub>) are divalent linking groups independently selected from the  
 5 group consisting of: a bond, -(CH<sub>2</sub>)<sub>m</sub>-C(OH)-, -(CH<sub>2</sub>)<sub>m</sub>-O-, -(CH<sub>2</sub>)<sub>m</sub>-S-, -(CH<sub>2</sub>)<sub>m</sub>-S(O)-, -(CH<sub>2</sub>)<sub>m</sub>-S(O)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>m</sub>-N(R<sub>40</sub>)-, -(CH<sub>2</sub>)<sub>m</sub>-C(R<sub>40</sub>)(R<sub>41</sub>)-, -(CH<sub>2</sub>)<sub>m</sub>-C(O)-, -N(R<sub>40</sub>)-C(O)-, -(CH<sub>2</sub>)<sub>m</sub>-CH=CH-, and -(CH<sub>2</sub>)<sub>m</sub>-C≡C-;

where m is 0-5;

R<sub>40</sub> and R<sub>41</sub> each is independently selected from: hydrogen, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-  
 10 C<sub>5</sub> hydroxyalkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, C<sub>3</sub>-C<sub>5</sub> cycloalkenyl;

Z<sub>P</sub> is selected from: branched C<sub>3</sub>-C<sub>5</sub> alkyl, C<sub>3</sub>-C<sub>10</sub> hydroxyalkyl, C<sub>3</sub>-C<sub>10</sub> hydroxyalkenyl, C<sub>3</sub>-C<sub>10</sub> hydroxyalkynyl, C<sub>3</sub>-C<sub>10</sub> hydroxycycloalkyl, C<sub>4</sub>-C<sub>10</sub> hydroxy cycloalkenyl, and oxocycloalkyl;

15 Z<sub>NP</sub> is selected from: C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, C<sub>3</sub>-C<sub>5</sub> cycloalkenyl, C<sub>1</sub>-C<sub>5</sub> hydroxyalkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, C<sub>1</sub>-C<sub>5</sub> alkylaryl, C<sub>1</sub>-C<sub>5</sub> hydroxyalkylaryl, C<sub>0</sub>-C<sub>5</sub> alkyl-CO<sub>2</sub>H, C<sub>0</sub>-C<sub>3</sub> alkyl-cycloalkyl-CO<sub>2</sub>H, C<sub>0</sub>-C<sub>5</sub> alkyl-N(R<sub>40</sub>)(R<sub>41</sub>), -X-(C<sub>1</sub>-C<sub>5</sub> alkyl), -X-(C<sub>1</sub>-C<sub>5</sub> alkenyl), -X-(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), -X-(C<sub>3</sub>-C<sub>5</sub> cycloalkenyl), -X-(C<sub>1</sub>-C<sub>5</sub> haloalkyl), -X-(C<sub>1</sub>-C<sub>5</sub> hydroxyalkyl), -X-(C<sub>1</sub>-C<sub>5</sub> alkylaryl), -X(OC<sub>1</sub>-C<sub>5</sub> alkyl), -XN(R<sub>40</sub>)(R<sub>41</sub>), -XN(R<sub>40</sub>)aryl, -N(CH<sub>3</sub>)(OCH<sub>3</sub>),  
 20 -N(OH)(CH<sub>3</sub>), -N(R<sub>42</sub>)-(C<sub>1</sub>-C<sub>5</sub> alkyl)CO<sub>2</sub>H, -N(R<sub>42</sub>)-(C<sub>1</sub>-C<sub>5</sub> alkyl)C(O)(C<sub>1</sub>-C<sub>5</sub> alkyl), -N(R<sub>42</sub>)-(C<sub>1</sub>-C<sub>5</sub> alkyl)C(O)(OC<sub>1</sub>-C<sub>5</sub> alkyl), -N(R<sub>42</sub>)-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>5</sub> alkyl), -NR<sub>42</sub>-S(O)-(C<sub>1</sub>-C<sub>5</sub> alkyl), -P(O)-(OC<sub>1</sub>-C<sub>5</sub> alkyl)<sub>2</sub>, heteroaryl, and -N=C(R<sub>40</sub>)N(R<sub>40</sub>)(R<sub>41</sub>);

25 R<sub>42</sub> is selected from: H, C<sub>1</sub>-C<sub>3</sub> alkyl, and C<sub>1</sub>-C<sub>3</sub> haloalkyl; and

X is selected from: O, C(O), C(S), S(O), and SO<sub>2</sub>;

provided that -(L<sub>NP</sub>)-Z<sub>NP</sub> is substituted at either the 12 or 13 position of the naphthalene ring; or

a pharmaceutically acceptable salt, solvate, prodrug, enantiomer, racemate,  
 30 diastereomer or mixture of diastereomers thereof.

-12-

It will be understood by those skilled in the art that the individual groups listed herein for the divalent linkers, (L<sub>P1</sub>), (L<sub>P2</sub>), and (L<sub>NP</sub>), can be attached at either end to the benzoxazole nucleus. For example, for the linker, -N(R<sub>40</sub>)-C(O)-, either the nitrogen can be attached to the naphthalene nucleus or alternatively the carbonyl carbon can be  
 5 attached to the naphthalene nucleus.

In preferred embodiments, compounds of the invention include the compound of Formula I having as preferred substituents;

R and R' are independently methyl or ethyl;

RP is hydrogen or methyl;

10 RP<sub>3</sub> and RN are independently hydrogen, methyl, ethyl, -O-methyl, or cyclopropyl;

(L<sub>P1</sub>) is a bond;

(L<sub>P2</sub>) is a bond, -CH<sub>2</sub>-, -CH(OH)-, or -C(Me)OH-;

(L<sub>NP</sub>) is a bond, -C(O)-, -C(O)NH-, or -C(O)N(Me)-;


15 Z<sub>P</sub> is 1,1-dimethylethyl, 1-hydroxycyclopentyl, 1-hydroxycyclohexyl, 3-ethyl-3-hydroxypentyl, 3-ethyl-3-hydroxypentenyl, or 3-ethyl-3-hydroxypentynyl,

Z<sub>NP</sub> is -CO<sub>2</sub>H, -CO<sub>2</sub>(R<sub>40</sub>), -N(R<sub>40</sub>)(R<sub>41</sub>), -NH-C(Me)(OH)-C(O)OH, -C(O)NMe-CH<sub>2</sub>-C(O)OH, -C(O)NMe-CH<sub>2</sub>-C(O)OMe, -C(O)NMe-CH<sub>2</sub>-C(O)OEt, -C(O)NMe-CH<sub>2</sub>-C(O)OiPr, -C(O)NMe-CH<sub>2</sub>-C(O)tBu, -cyclopropyl-C(O)OH, 20 -cyclobutyl-C(O)OH, -C(O)NMe-C(Me)<sub>2</sub>-C(O)OH, -C(O)N(R<sub>40</sub>)S(O)(R<sub>42</sub>), -C(O)N(R<sub>40</sub>)SO<sub>2</sub>R<sub>42</sub>, -C(O)-N(R<sub>40</sub>)-5-tetrazolyl, -C(O)N(R<sub>40</sub>)-(C<sub>1</sub>-C<sub>5</sub> alkyl)-S(O)R<sub>42</sub>, -C(O)N(R<sub>40</sub>)-(C<sub>1</sub>-C<sub>5</sub> alkyl)-S(O)<sub>2</sub>R<sub>42</sub>, C<sub>0</sub>-C<sub>3</sub> alkyl(cycloalkyl)C(O)NHSO<sub>2</sub>Me, C<sub>0</sub>-C<sub>3</sub> alkyl(cycloalkyl)-C(O)OH, C<sub>0</sub>-C<sub>3</sub> alkyl(cycloalkyl)-C(O)NH-heteroaryl, C<sub>0</sub>-C<sub>3</sub> alkyl(cycloalkyl)-NHSO<sub>2</sub>(C<sub>1</sub>-C<sub>5</sub> alkyl), 25 and -CH<sub>2</sub>CO<sub>2</sub>H.


Preferred compounds of the invention and salts and prodrug derivatives are represented by formulae C1 to C8 as follows:

-13-

C1)




C2)




5


C3)



C4)




C5)




10

-14-

C6)




C7)



5

C8)



where R1 is H, Me, or Et; R2 is H or Me; R3 is H Me, or Et; and R4 is H or Me.

Particularly preferred compounds include compounds represented by formulae C1-C8

10 where R1 is a Me or Et; and R2 and R3 individually are H or Me.

### EXAMPLES

#### General Experimental Conditions:

15 The starting material/intermediate is the compound from the immediate preceding experimental unless otherwise indicated.

All reactions are performed under nitrogen/argon atmosphere, in a stirred reaction vessel, and at room temperature unless indicated otherwise.

Unless otherwise indicated, the notation that "organic layer is MgSO<sub>4</sub>/Na<sub>2</sub>SO<sub>4</sub> dried" "dried over MgSO<sub>4</sub>/Na<sub>2</sub>SO<sub>4</sub>" is defined as swirling or stirring the solution with a

-15-

dessicant ( $\text{MgSO}_4$  and/or  $\text{Na}_2\text{SO}_4$ ) 5-15 m, then filtering off the dessicant to give an anhydrous filtrate.

For analogous multi-step reaction procedures, the yield is given either for the ultimate step or overall multi-steps as indicated.

5 Solutions are "concentrated" at a range of 25-75 °C with reduced pressure (0.05 to 1 mm).

Unless otherwise indicated, "the residue is chromatographed" is defined as silica gel chromatography of residue with moderate nitrogen pressure (flash chromatography) or a medium pressure chromatography systems using a silica gel to crude product ratio of  
10 ~10-100.

For HPLC, the conditions listed are for the analytical trace only. For Preparative HPLC, the eluent is similar to analytical HPLC eluent.

Thin layer chromatography is performed with silica gel plates with UV and/or appropriate staining solution.

15 NMR spectra are obtained with either 300 or 400 MHz spectrometer. NMR data is listed to denote spectrum is consistent with assigned structure. "NMR" notation without data denotes spectrum is consistent with assigned structure.

HRMS – high resolution mass spectrum

ES-MS – electrospray mass spectrum

20

Abbreviations:

Aq - aqueous

d – day

eq - equivalent

25

h – hour

m – minute

satd – saturated

disp – dispersion

quant – quantitative

30

rt for retention time (both small caps to minimize confusion with RT)

RT – room temperature



-16-

TABLE 1  
Chemical Terms

Term	Definition	Term	Definition
BF <sub>3</sub> -OEt <sub>2</sub>	boron trifluoride etherate	MeOH	methanol
BnBr	benzyl bromide	NMO	4-methylmorpholine N-oxide
CH <sub>2</sub> Cl <sub>2</sub>	Dichloromethane	NMP	N-methylpyrrolidin- 2-one
DMAP	4-(dimethylamino)pyridine	Na-S-R <sub>3</sub>	sodium alkylmercaptide
DMF	N,N-dimethylformamide	PBr <sub>3</sub>	phosphorus tribromide
DMSO	Dimethylsulfoxide	Pd(DPPF)	palladium dichloro[1,1'- bis(diphenylphosphi no)ferrocene
DPPB	1,4-bis(diphenylphosphino) butane	Pd(OAc) <sub>2</sub>	palladium (II) acetate
DPPF	dichloro[1,1'- bis(diphenylphosphino) ferrocene	Pd(TPP) <sub>4</sub>	palladium tetrakis(triphenylphos phine
EDCI	3-Ethyl-1-[3-(dimethylamino) propyl]carbodiimide hydrochloride	Pd-C	palladium on carbon
EEDC	Diethyl cyanamide	Pd-C/H <sub>2</sub>	palladium on carbon

-17-

Term	Definition	Term	Definition
			with hydrogen pressure
EtMgBr	ethyl magnesium bromide	pTSA	para-toluenesulfonic acid
EtOAc	ethyl acetate	Pyr	pyridine
EtOH	Ethanol	Red-Al	sodium bis(2-methoxyethoxy)aluminum hydride
H <sub>2</sub>	hydrogen pressure	R <sub>2</sub> MgBr	alkyl magnesium bromide
H <sub>2</sub> NCH <sub>2</sub> CO 2Me	methyl glycinate	R <sub>3</sub> MgBr	alkyl magnesium bromide
Hept	Heptane	R <sub>5</sub> MgBr	alkyl magnesium bromide
Hex	Hexanes	R <sub>3</sub> S(O) <sub>2</sub> Cl	alkylsulfonyl chloride
HN(OMe)Me	N-methyl-O-methyl hydroxylamine	R <sub>2</sub> S(O) <sub>2</sub> N	alkylsulfonamide
HNMe <sub>2</sub>	dimethyl amine	H <sub>2</sub>	
HATU	O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate	TBSCl	<i>tert</i> -butyldimethylsilyl chloride
HOAT	7-aza-1-hydroxy benzotriazole	tBuC(O)C H <sub>2</sub> Br –	1-bromopinacolone
		Tf <sub>2</sub> O	triflic anhydride

-18-


Term	Definition	Term	Definition
HOBT	1-hydroxybenzotriazole	TFA	trifluoroacetic acid
K <sub>2</sub> CO <sub>3</sub>	potassium carbonate	THF	tetrahydrofuran
LAH	Lithium aluminum hydride	Ti(OiPr) <sub>4</sub>	titanium tetraisopropoxide
LDA	Lithium diisopropyl amide	TMS-	trimethylsilyl
LiHMDS	Lithium hexamethyl disilazide	acetylene	acetylene
Lindlar catalyst	Pd-CaCO <sub>3</sub> -PbO	TPAP	tetrapropylammoniu m perruthenate
mCPBA	meta-chloroperbenzoic acid	BnHalide	benzylhalide
TPA	12- <i>O</i> -tetradecanoyl 13-acetate (Sigma)	PHA	Phytohemagglutinin (Sigma)
NaHMDS	Sodium hexamethyl disilazide		
NMM	N-methylmorpholine		

-19-

## General Procedures

## Scheme

1



-20-

6-Hydroxy-2-naphthoic acid is esterified in an alkanol with acid catalysis (HCl, sulfuric acid, or toluenesulfonic acid) from room temperature to the reflux temperature of the mixture to give the ester 1. Ester 1 is protected, e.g., with a benzylhalide and base to give the protected ester 2. Ester 2 is treated with excess alkyl

5 Grignard reagent (2 to 5 equivalents) in diethylether or THF from 0 ° C. to room temperature to give the tert-carbinol 3. Carbinol 3 is hydrogenated with a catalyst, e.g., palladium on carbon to give the deprotected carbinol 4. Carbinol 4 is reacted with triflic anhydride to produce the triflate that dehydrates upon work up to the Z/E-olefin 5. Triflate 5 is reacted with carbon monoxide (1 to 100 psi) and a palladium


10 catalyst (0.1 to 10%) in alkanol with DMF from room temperature to 150 ° C. for 8 to 48 h to give the ester 6. Ester 6 alkylates an ortho-substituted phenol in the presence of a Lewis acid, e.g., boron trifluoride etherate (0.01 to 5 equivalents) at room temperature to give the diarylmethane 7. A variety ortho-substitued phenols are commercially available or can be readily prepared by those skilled in the art. The free

15 hydroxyl of the diarylmethane 7 is alkylated with an alpha-halo ketone (z-C(O)CH<sub>2</sub>hal, where z is an alkyl group or a substituted alkyl group) to give keto ester 8. Keto ester 8 is saponified with lithium, sodium, or potassium hydroxide in alkanol from room temperature to the reflux temperature of the mixture to give the keto acid 9.

20 Keto acid 9 is reduce to the carbinol 10. with lithium or sodium borohydride, or cyanoborohydride in alkanol at room temperature.

-21-

Scheme II.




## Scheme II.

2,7-Dihydroxynaphthalene is reacted with triflic anhydride (0.9 to 1.3 equivalents) to produce the mono-triflate 11. Triflate 11 is reacted with carbon monoxide (1-100 psi) in DMF with alkanol and a palladium catalyst from room temperature to 150 ° C. and from 8 to 48 h to give the ester 12. The ester 12 is protected, e.g., with a benzylhalide and base to give the protected ester 13. Ester 13 is reacted with an excess of alkyl Grignard reagent (2-5 equivalents) to give the tert-carbinol 14. The protection group of 14 is removed, e.g., by hydrogenation in alkanol with a palladium catalyst to give the unprotected tert-carbinol 15. Carbinol 15 is converted to triflate 16. with triflic anhydride and base. Triflate 16 is converted to the ester 17 as above with carbon monoxide in DMF/alkanol and base in the presence of a palladium catalyst, e.g., palladium acetate and DPPB. Ester 17 alkylates an ortho-substituted phenol in the presence of a Lewis acid, e.g., boron trifluoride etherate (0.01 to 5 equivalents) at room temperature to give the diarylmethane 18. The free hydroxyl of the diarylmethane 18 is alkylated with an alpha-halo ketone( $\text{z-C(O)CH}_2\text{hal}$ , where z is an alkyl group or a substituted alkyl group) to give keto ester 19. Keto ester 19 is saponified with lithium, sodium, or potassium hydroxide in alkanol from room temperature to the reflux temperature of the mixture to give the keto acid 20. Reduction of keto acid 20 to the carbinol is achieved as above with lithium or sodium borohydride or cyanoborohydride in alkanol at room temperature.

Each of the free acids produced in each of the above schemes (9, 10, and 20) are converted to esters and carboxamides using reactions well known to those skilled in the art.

-23-

## Scheme III




## Scheme III.

- 5 Alcohol 17 is reacted with  $\text{BF}_3\cdot\text{OEt}_2$  and o-cresol (or another ortho substituted phenol) to give phenol 21. Phenol 21 is reacted with an alpha haloketone ( $\text{z-C(=O)CH}_2\text{hal}$ , where z is an alkyl group or a substituted alkyl group) and a base to give ketone 22. Ketone 22 is treated with a hindered base (LDA or NaHMDS) and either methyl halide or ethyl halide (or other alkyl halide) to afford ketone 23. Ketone
- 10 23 is reacted with an alkali hydroxide to give acid 24. Acid 24 is coupled with substituted amino acid ester using EDCI/HOBT/NMM (N-methylmorpholine) to amide-ester 26. Coupling of acid 24 with cyclic amino acid ester the corresponding cyclic amide-ester 26 is obtained. Amide-ester 26 is reacted with alkali hydroxide to give amide 27.



-24-

## Scheme IV




-25-

## Scheme IV.


Ketone-acid 24 is reacted with NaBH<sub>4</sub>/MeOH to give alcohol-acid 28. Alcohol-acid 28 is coupled with substituted sulfonamide using EDCI/DMAP to afford acylsulfonamide 29. Alcohol-acid 28 is reacted with EDCI/5-aminotetrazole/DMAP to give acylaminotetrazole 30. (The 5-aminotetrazole can be replaced with other heterocyclics or heteroaryl groups as desired.) Alcohol-acid 28 is reacted with formamide and NaOMe at 100 °C to produce amide 31. Amide 31 is reacted with Et<sub>3</sub>N, 2-chloro-1,3-dimethylimidazolinium hexafluorophosphate, and TFA to afford nitrile 32. Nitrile 32 is reacted with Bu<sub>3</sub>SnN<sub>3</sub> at 80 °C to give tetrazole 33.

10

Example 1. Preparation of 6-{1-[4-(3,3-dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid




15 A. 6-Hydroxynaphthalene-2-carboxylic acid methyl ester



Treat a mixture of 6-hydroxy-2-naphthoic acid (4.45g, 23.6 mmol) in 2,2-dimethoxypropane (235 mL) with conc. HCl (24 mL) and then MeOH (60 mL). Stir the reaction at RT for 16 h and then at 55 °C for 16 h. Pour the reaction mixture into EtOAc (250 mL) and wash with brine (3 x 100 mL). Dry the organic layer over Na<sub>2</sub>SO<sub>4</sub>, concentrate the filtrate, and chromatograph the concentrated filtrate (0.5 kg silica gel, 5:95 to 15:85 EtOAc:hex) to give the title compound (4.77 g, quant). MS (ES) *m/e* 203 (M+1).


-26-

## B. 6-Benzyloxynaphthalene-2-carboxylic acid methyl ester



Treat 6-hydroxynaphthalene-2-carboxylic acid methyl ester (4.77 g, 23.6 mmol) and benzyl bromide (3.3 mL, 28.3 mmol) in DMF (22 mL) with cesium carbonate (15.3 g, 47.2 mmol). Stir the mixture for 22 h at RT and concentrate. Dilute the concentrate with EtOAc (300 mL) and water (100 mL). Wash the organic layer with brine (3 x 100 mL), dry it over Na<sub>2</sub>SO<sub>4</sub>, and filter. Concentrate the filtrate. Purify the product using medium pressure silica gel chromatography (15:85 to 50:50 EtOAc:hex) to give the title compound as a white solid (5.75 g, 83%). MS (ES) *m/e* 293 (M+1).


## C. 3-(6-Benzyloxynaphthalen-2-yl)pentan-3-ol



Cool 6-benzyloxynaphthalene-2-carboxylic acid methyl ester (5.75 g, 19.67 mmol) in THF (115 mL) to about 0 °C and treat the solution dropwise with ethyl magnesium bromide (23.0 mL, 69.0 mmol, 3.0 M in ether). Allow the reaction to warm to RT and stir it for 3 h. Quench with water and concentrate. Dissolve the crude residue in CH<sub>2</sub>CL<sub>2</sub> (200 mL), wash twice with brine, dry over Na<sub>2</sub>SO<sub>4</sub> and filter. Concentrate the filtrate to give the title compound as a yellow solid (6.0 g, 95%). MS (ES) *m/e* 321 (M+1).


## D. 6-(1-Ethyl-1-hydroxypropyl)naphthalen-2-ol

-27-




Treat a mixture of 3-(6-benzyloxynaphthalen-2-yl)pentan-3-ol (1.5 g, 4.68 mmol) and 5% Pd/Al<sub>2</sub>CO<sub>3</sub> (0.90 g) in ethanol (750 mL) with hydrogen at 60 psi for 8 h at RT. Filter the catalyst from the reaction mixture and concentrate the filtrate to  
 5 give the title compound as a yellow solid (0.94 g, 87%). MS (ES) *m/e* 229 (M-1).

E. Trifluoromethanesulfonic acid 6-(1-ethylpropenyl)naphthalen-2-yl ester



Cool a solution of 6-(1-ethyl-1-hydroxypropyl)naphthalen-2-ol (0.94 g, 4.08 mmol) and pyridine (1.3 mL, 16.3 mmol) in CH<sub>2</sub>CL<sub>2</sub> (30 mL) in an ice bath and treat  
 10 dropwise with trifluoromethanesulfonic anhydride (1.0 mL, 6.1 mmol). Remove the cooling bath, and stir the mixture at RT for 1h. Quench the reaction mixture with ice water. Dilute the mixture with CH<sub>2</sub>CL<sub>2</sub> (80 mL), wash twice with brine, dry over Na<sub>2</sub>SO<sub>4</sub> and filter. Concentrate the filtrate and purify by silica gel radial  
 15 chromatography (CH<sub>2</sub>CL<sub>2</sub>) to give the title compound as a yellow oil (0.93 g, 66%). MS (ES) *m/e* 361 (M+NH<sub>4</sub>).

F. 6-(1-Ethylpropenyl)naphthalene-2-carboxylic acid methyl ester




20 Treat a mixture containing trifluoromethanesulfonic acid 6-(1-ethylpropenyl)naphthalen-2-yl ester (0.92 g, 2.67 mmol), Pd(OAc)<sub>2</sub> (0.020 mg, 0.09

-28-

mmol), *bis*-(diphenylphosphino)ferrocene (0.046 g, 0.083 mmol), Et<sub>3</sub>N (0.67 mL, 4.81 mmol), in MeOH (10 mL) and DMSO (15 mL) with carbon monoxide at 100 psi at 80 °C for 4 h. Pour the reaction mixture into ether (100 mL) and wash the mixture with brine (4 x 50 mL). Dilute the organic layer with EtOAc (100 mL), wash with  
5 brine, then dry over Na<sub>2</sub>SO<sub>4</sub> and filter. Concentrate the filtrate to give the title compound as an off-white solid (0.67 g, quant). MS (ES) *m/e* 255 (M+1).


G. 6-[1-Ethyl-1-(4-hydroxy-3-methylphenyl)propyl]naphthalene-2-carboxylic acid methyl ester



10

Cool a solution of 6-(1-ethyl-propenyl)naphthalene-2-carboxylic acid methyl ester (0.74 g, 2.75 mmol) and *o*-cresol (2.0 mL, 19.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) to -78 °C; dropwise add BF<sub>3</sub>·OEt<sub>2</sub> (1.22 mL, 9.6 mmol). Remove the cooling bath and stir the reaction at RT for 16 h. Quench the reaction with ice water; then dilute with  
15 EtOAc (200 mL). Wash the resulting mixture twice with brine, dry over Na<sub>2</sub>SO<sub>4</sub>, filter, and concentrate the filtrate. Purify the product using medium pressure silica gel chromatography (100% hex to 25:75 EtOAc:hex) to give the title compound as a white solid (0.87 g, 87%). MS (ES) *m/e* 363 (M+1).

20 H. 6-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid methyl ester




Stir a mixture containing 6-[1-ethyl-1-(4-hydroxy-3-methylphenyl)propyl]naphthalene-2-carboxylic acid methyl ester (0.87 g, 2.40 mmol),

-29-

1-bromopinacolone (0.64 g, 3.60 mmol), and potassium carbonate (1.0 g, 7.2 mmol) in acetone (15 mL) at RT for 16 h. Concentrate the reaction mixture and dilute the residue with EtOAc (100 mL) and water (50 mL). Adjust the pH of the mixture to 1 using 1 N HCl. Discard the aqueous layer and wash the organic layer with brine, dry over Na<sub>2</sub>SO<sub>4</sub>, then filter. Concentrate the filtrate and purify by silica gel radial chromatography (5:95 EtOAc:hex to 15:85 EtOAc:hex) to give the title compound as an off-white solid (0.53 g, 48%). MS (ES) *m/e* 461 (M+1).


10 1. 6-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid



Treat a solution of 6-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}-naphthalene-2-carboxylic acid methyl ester (0.53 g, 1.15 mmol) in THF (8 mL) and MeOH (16 mL) with 2N NaOH (2.9 mL, 5.8 mmol) and heat at 55 °C for 16 h. Concentrate the reaction mixture. Dilute the residue with in EtOAc (50 mL) and water (25 mL) and acidify the resulting mixture to pH 1 using 1N HCl. Discard the aqueous layer and wash the organic layer with brine, Dry the organic layer over Na<sub>2</sub>SO<sub>4</sub>, filter, and concentrate the filtrate to give the title compound as a yellow solid (0.45 g, 87%). MS (ES) *m/e* 447 (M+1).

20


Example 2. Preparation of 6-{1-Ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylic acid



-30-

Treat a solution of 6-{1-[4-(3,3-dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid (0.15 g, 0.34 mmol) in THF (6 mL) with NaBH<sub>4</sub> (0.025 g, 0.67 mmol) and stir the reaction mixture at RT for 1 h. Quench the reaction first with water (5 mL) and then with 1N HCl (2 mL). Remove the THF under vacuum. Dilute the mixture with EtOAc (50 mL), wash with brine (2x), dry over NaSO<sub>4</sub> and filter. Concentrate the filtrate to give the title compound (0.15 g). MS (ES) *m/e* 447 (M-1).

Example 2A and 2B. Preparation of enantiomers of 6-{1-Ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylic acid



Chromatograph a mixture of racemic 6-{1-ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylic acid (2.93 g) on a Chiralpak AD-H column, Daicel Chemical Industries, to give enantiomer 1, Example 2A (1.42 g, 48%) and enantiomer 2, Example 2B (1.38 g, 47%).

Enantiomer 1, Example 2A.

HPLC: ChiralPak AD-H (4.6x150); 60%heptane/40%IPA/0.1%TFA; 0.6 mL/m (flow rate); uv: 250 nm.

rt = 10 m


Enantiomer 2, Example 2B

HPLC: ChiralPak AD-H (4.6x150); 60%heptane/40%IPA/0.1%TFA; 0.60 mL/m (flow rate); uv: 250 nm.

rt = 16 m


-31-

Example 3. Preparation of [(6-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carbonyl)amino]acetic acid



Treat a solution of 6-{1-[4-(3,3-dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid (0.11 g, 0.246 mmol) in CH<sub>2</sub>CL<sub>2</sub> (1.0 mL) dropwise with oxalyl chloride (0.13 mL, 1.48 mmol). Stir the solution at RT for 1 h and concentrate to give 6-{1-[4-(3,3-dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carbonyl chloride ("the acid chloride"). Treat a mixture of methylglycine HCl (0.031 g, 0.25 mmol) and Et<sub>3</sub>N (0.51 mL, 0.37 mmol) in CH<sub>2</sub>CL<sub>2</sub> (3 mL) with a solution of the acid chloride (0.057 g, 0.123 mmol) in CH<sub>2</sub>CL<sub>2</sub> (1 mL). Stir the reaction at RT for 16 h and concentrate. Dissolve the residue in THF (0.5 mL) and MeOH (1.0 mL) and treat with 2N NaOH (0.24 mL, 0.48 mmol). Heat the solution to 50 °C for 16 h, concentrate, dilute with CH<sub>2</sub>CL<sub>2</sub> and water (10 mL each). Acidify the resulting mixture to pH 1 using 5N HCl. Wash the organic layer with brine, dry over Na<sub>2</sub>SO<sub>4</sub>, and filter. Concentrate the filtrate and purify by silica gel radial chromatography (10:90 MeOH:CH<sub>2</sub>CL<sub>2</sub>) to give the title compound as a white solid (0.021 g, 34%). MS (ES) *m/e* 504 (M+1).

Example 4. Preparation of 6-{1-[4-(3,3-dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid dimethylamide




Treat a solution of dimethylamine (2.0 M in THF) (0.12 mL, 0.25 mmol) and




-32-

Et<sub>3</sub>N (0.034 mL, 0.25 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) with 6-{1-[4-(3,3-dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carbonyl chloride (0.057 g, 0.123 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL). Stir the mixture at RT for 16 h; dilute with CH<sub>2</sub>Cl<sub>2</sub> and water. Acidify the resulting mixture to pH 1 using 5N HCl. Separate the organic layer from the aqueous layer. Wash the organic layer with brine, dry over Na<sub>2</sub>SO<sub>4</sub>, and filter. Concentrate the filtrate to give the title compound as a yellow solid (0.033 g, 57%). MS (ES) *m/e* 474 (M+1).

Example 5. Preparation of [(6-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carbonyl)methylamino]acetic acid



A. [(6-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carbonyl)methylamino]acetic acid ethyl ester




Dissolve 6-{1-[4-(3,3-dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid (0.10 g, 0.222 mmol), sarcosine ethyl ester HCl (0.069 g, 0.45 mmol), DMAP (0.108 g, 0.89 mmol) and EEDC (0.086 g, 0.45 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL). Stir the mixture at RT for 16 h. Dilute the reaction mixture with CH<sub>2</sub>Cl<sub>2</sub> and water. Acidify to pH 1 using 1N HCl. Separate and discard the aqueous layer. Wash the organic layer with brine, dry over Na<sub>2</sub>SO<sub>4</sub>, and filter. Concentrate the filtrate and purify by silica gel radial chromatography (25:75 EtOAc:hex to 50:50 EtOAc:hex) to give the title compound as a white solid (0.093 g, 78%). MS (ES) *m/e* 546 (M+1).

-33-


B. [(6-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carbonyl)methylamino]acetic acid

Treat a solution of [(6-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carbonyl)methylamino]acetic acid ethyl ester (0.090 g, 0.16 mmol) in THF (1.25 mL) and EtOH (2.5 mL) with 2N NaOH (0.80 mL, 1.6 mmol). Heat the mixture to 50 °C for 16 h. Acidify the reaction mixture to pH 1 using 1N HCl. Concentrate the resulting mixture. Dilute the residue with EtOAc and water (25 mL each). Separate and discard the aqueous layer. Wash the organic layer with brine, dry over Na<sub>2</sub>SO<sub>4</sub>, and filter. Concentrated the filtrate to give the title compound as a white solid (0.070 g, 85%). MS (ES) *m/e* 518 (M+1).

Example 6. Preparation of 7-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid



A. Trifluoromethanesulfonic acid 7-(1-ethyl-1-hydroxypropyl)naphthalen-2-yl ester




Cool a solution of 2,7-dihydroxynaphthalene (2.7 g, 31.2 mmol) and pyridine (4.53 mL, 56.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) in an ice bath and treat the cooled solution with trifluoromethanesulfonic anhydride (6.62 mL, 39.3 mmol). Remove the cooling bath and stir the mixture at RT for 2.5 h. Quench the reaction with ice water and dilute with CH<sub>2</sub>Cl<sub>2</sub> (200 mL). Wash the residue with water (2 x 100 mL) and then acidify to pH 3 using 1N HCl. Dry the organic layer over Na<sub>2</sub>SO<sub>4</sub>, filter and concentrate the filtrate. Purify the product by medium pressure silica gel

-34-


chromatography (10:90 EtOAc:hex to 25:75 EtOAc:hex) to give the title compound as a yellow oil (4.1 g, 45%). MS (ES)  $m/e$  291 (M+1).

B. 7-Hydroxynaphthalene-2-carboxylic acid methyl ester



5 Treat a mixture containing trifluoromethanesulfonic acid 7-(1-ethyl-1-hydroxypropyl)naphthalen-2-yl ester (0.41 g, 14.1 mmol), Pd(OAc)<sub>2</sub> (0.30 g, 1.38 mmol), *bis*-(diphenylphosphino)ferrocene (0.705 g, 1.27 mmol), Et<sub>3</sub>N (10.2 mL, 73.2 mmol), in MeOH (12 mL) and DMSO (18 mL) with carbon monoxide at 100 psi at 80  
10 °C for 4 h. Pour the reaction mixture into ether (300 mL) and wash with brine (5 x 100 mL). Add EtOAc (250 mL), wash the organic layer with brine, dry over Na<sub>2</sub>SO<sub>4</sub>, and filter. Concentrate the filtrate to give the title compound as a tan solid (2.84 g, quant). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 3.98 (s, 3H), 7.23-7.29 (m, 2H), 7.78 (t, J= 8.3 Hz, 2H), 7.88-7.90 (m, 1H), 8.43 (d, J= 0.88 Hz, 1H).


15 C. 7-Benzoyloxynaphthalene-2-carboxylic acid methyl ester



Treat a solution of 7-hydroxynaphthalene-2-carboxylic acid methyl ester (2.84 g, 14.0 mmol) and benzyl bromide (1.84 mL, 15.5 mmol in DMF (14 mL) with  
20 cesium carbonate (9.10 g, 27.9 mmol). Stir the mixture for 2 h and concentrate. Dissolve the crude residue in EtOAc (150 mL) and 1 N HCl (50 mL). Wash the organic layer with brine (3 x 100 mL), dry over Na<sub>2</sub>SO<sub>4</sub>, and filter. Concentrate the filtrate and purify using medium pressure silica gel chromatography (10:90 EtOAc:hex) to give the title compound as a white solid (1.47 g, 36%). MS (ES)  $m/e$   
25 293 (M+1).


-35-

## D. 3-(7-Benzyloxynaphthalen-2-yl)pentan-3-ol



Cool a THF (30 mL) solution of 7-benzyloxynaphthalene-2-carboxylic acid methyl ester (1.47 g, 5.03 mmol) with an ice bath and treat the cooled solution dropwise with ethyl magnesium bromide (5.9 mL, 17.6 mmol, 3.0M in ether). Remove the ice bath and stir the reaction for 3 h. Quench the reaction with ice water, then treat with 1 N HCl, and concentrate to provide a crude product. Dissolve the crude product in EtOAc (100 mL). Wash the organic layer twice with brine, dry over Na2SO4, and filter. Concentrate the filtrate and purify using radial silica gel chromatography (5:95 to 15:85 EtOAc:hex) to give a yellow solid (0.46 g, 90%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 0.787 (t, J= 7.5 Hz, 6H), 1.67 (br s, 1H), 1.82-2.02 (m, 4H), 5.19 (s, 2H), 7.19-7.44 (m, 6H), 7.50 (d, J= 6.6 Hz, 2H), 7.73 (d, J= 8.8 Hz, 2H), 7.77 (d, J= 1.8 Hz, 1H).


## E. 7-(1-Ethyl-1-hydroxypropyl)naphthalen-2-ol



Treat a mixture of 3-(7-benzyloxynaphthalen-2-yl)pentan-3-ol (1.45 g, 4.53 mmol) and 5% Pd/Al<sub>2</sub>CO<sub>3</sub> (0.044 g) in EtOH (725 mL) with hydrogen at 60 psi for 8 h at RT. Remove the catalyst by filtration and concentrate the filtrate to give the title compound as a yellow oil (0.98 g, 94%). MS (ES) *m/e* 229 (M-1).


-36-

## F. Trifluoromethanesulfonic acid 7-(1-ethyl-1-hydroxypropyl)naphthalen-2-yl ester



Cool a CH<sub>2</sub>CL<sub>2</sub> (30 mL) solution of 7-(1-ethyl-11-hydroxypropyl)naphthalen-2-ol (0.97 g, 4.2 mmol) and pyridine (1.3 mL, 16.3 mmol) with an ice bath. Treat the cooled solution dropwise with trifluoromethanesulfonic anhydride (1.0 mL, 6.1 mmol). Remove the cooling bath and stir the mixture for 1.5 h at RT. Quench the reaction with ice water. Dilute the reaction mixture with EtOAc and water (100 mL each) and acidify to pH 1 using 0.1N HCl. Dry the organic layer over Na<sub>2</sub>SO<sub>4</sub>, filter and concentrate the filtrate. Purify by radial silica gel (2:98 to 10:90 EtOAc:hex) to give the title compound as a colorless oil (1.46 g, 66%). MS (ES) *m/e* 377 (M-NH<sub>4</sub>).


## G. 7-(1-Ethyl-1-hydroxypropyl)naphthalene-2-carboxylic acid methyl ester



Treat a mixture of trifluoromethanesulfonic acid 7-(1-ethyl-1-hydroxypropyl)naphthalen-2-yl ester (1.0 g, 2.76 mmol), Pd(OAc)<sub>2</sub> (0.062 g, 0.27 mmol), *bis*-(diphenylphosphino)ferrocene (0.14 g, 0.25 mmol), Et<sub>3</sub>N (2.0 mL, 14.3 mmol) in MeOH (10 mL) and DMSO (15 mL) with carbon monoxide at 100 psi at 80 °C for 4 h. Pour the reaction mixture into ether (100 mL) and wash the resulting solution with brine (4 x 50 mL). Add EtOAc (100 mL), wash the organic layer with brine. Dry the organic layer over Na<sub>2</sub>SO<sub>4</sub>, filter, and concentrate the filtrate. Purify by silica gel radial chromatography (5:95 to 15:85 EtOAc:hex) to give the title compound as a yellow oil (0.71 g, 94%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 0.792 (t, J= 7.5 Hz, 6H), 1.80-2.04 (m, 4H), 4.00 (s, 3H), 7.58 (dd, J= 8.8, 1.4 Hz, 1H), 7.80-7.87 (m, 2H), 7.99 (d, J= 1.3 Hz, 1H), 8.02 (d, J= 1.8 Hz, 1H), 8.05 (d, J= 1.8 Hz, 1H), 8.61 (d, J= 1.0 Hz, 1H).


-37-

H. 7-[1-Ethyl-1-(4-hydroxy-3-methylphenyl)propyl]naphthalene-2-carboxylic acid methyl ester



Cool a CH<sub>2</sub>CL<sub>2</sub> (20 mL) solution of 7-(1-ethyl-1-hydroxypropyl)naphthalene-2-carboxylic acid methyl ester (0.73 g, 2.68 mmol) and *o*-cresol (1.54 mL, 15.0 mmol) to -78 °C; then dropwise add BF<sub>3</sub>-OEt<sub>2</sub> (0.76 mL, 6.0 mmol). Allow the reaction mixture to warm to RT and stir for 1 h. Quench the reaction with ice water. Dilute the resulting mixture with EtOAc (200 mL). Wash the mixture with brine, dry the organic layer over Na<sub>2</sub>SO<sub>4</sub>, filter and concentrate the filtrate. Purify by silica gel radial chromatography (5:95 to 35:65 EtOAc:hex) to give the title compound as a yellow oil (0.93 g, 86%). MS (ES) *m/e* 363 (M+1).

I. 7-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid methyl ester




Stirr a mixture of 7-[1-ethyl-1-(4-hydroxy-3-methylphenyl)propyl]naphthalene-2-carboxylic acid methyl ester (0.87 g, 2.40 mmol), 1-bromopinacolone (1.54 g, 8.68 mmol), and potassium carbonate (2.4 g, 17.4 mmol) in acetone (20 mL) for 4 h at RT. Concentrate the reaction mixture and then dilute the residue with EtOAc (100 mL) and water (50 mL). Adjust the pH of the mixture to 1 using 1 N HCl. Discard the aqueous layer. Wash the organic layer with brine, dry over Na<sub>2</sub>SO<sub>4</sub> and filter. Concentrate the filtrate and purified by silica gel radial chromatography (5:95 EtOAc:hex to 15:85 EtoAc:hex) to give the title compound as a white solid (1.1 g, 82%). MS (ES) *m/e* 461 (M+1).

25


-38-

J. 7-{1-[4-(3,3-Dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid



Treat 7-{1-[4-(3,3-dimethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid methyl ester (1.1 g, 2.4 mmol) in THF (15 mL) and MeOH (30 mL) with 2N NaOH (6.0 mL, 12.0 mmol) and heat to 55 °C for 16 h. Then concentrate the reaction mixture and dilute the residue with CH<sub>2</sub>CL<sub>2</sub> and water (50 mL each). Acidify the mixture to pH 1 using 1N HCl. Discard the aqueous layer and wash the organic layer with brine. Dry the organic layer over Na<sub>2</sub>SO<sub>4</sub>, filter and concentrate the filtrate. Purify by silica gel radial chromatography (2:98 MeOH:CH<sub>2</sub>CL<sub>2</sub> to 10:90 MeOH:CH<sub>2</sub>CL<sub>2</sub>) to give the title compound as a yellow solid (0.96 g, 90%). MS (ES) *m/e* 447 (M+1).

Example 7. Diastereomeric isomer pair 1 of 2-[6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl]-naphthalene-2-carboxylamino]acetic acid



A. 4-Bromo-2, 2-dimethylhexane-3-one


Slowly add bromine (10.87 mL, 212.15 mmol) to 2, 2-dimethylhexane-3-one (27.20 g, 212.15 mmol) in ether (200 mL) and allow the reaction to stir for 14 h. Combine the reaction mixture with water (200 mL) and partition. Dry the organic layer over Na<sub>2</sub>SO<sub>4</sub>, filter and then concentrate the filtrate to provide the title compound as a yellow oil (48.2 g, quant). <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 1.01 (t, J

-39-

= 7.6 Hz, 3H), 1.27 (s, 9H), 2.03 (m, 2H), 4.58 (t,  $J = 7.2$  Hz, 1H). High Res. ES-MS: 207.0348; calc. for  $C_8H_{15}BrO+H$ : 207.0384.

B. 6-{1-[4-(3,3-Dimethyl-1-ethyl-2-oxobutoxy)-3-methylphenyl]-1-

5 ethylpropyl}naphthalene-2-carboxylic acid methyl ester




Add 4-bromo-2, 2-dimethylhexane-3-one (660 mg, 3.187 mmol) to a mixture of 6-[1-ethyl-1-(4-hydroxy-3-methylphenyl)propyl]naphthalene-2-carboxylic acid methyl ester (1.1 g, 3.035 mmol), and  $K_2CO_3$  (629 mg, 4.55 mmol) in DMF (10 ml).

10 Maintain the reaction mixture  $45^\circ C$  for 14 h. Add an additional amount of 4-bromo-2, 2-dimethylhexane-3-one (314 mg, 1.52 mmol) and maintain the mixture at  $60^\circ C$  while stirring for 62 h. Cool the reaction mixture and dilute it with  $CH_2Cl_2$ , filter, and concentrate in vacuo. Chromatograph the resulting residue ( $CH_2Cl_2$  to 1% EtOAc/ $CH_2Cl_2$ ) to give the title compound as a colorless viscous oil (1.299 g, 88%).

15  $^1NMR$  (400MHz,  $CDCl_3$ )  $\delta$  ppm: 0.62 (t,  $J = 7.2$  Hz, 6H), 1.01 (t,  $J = 7.6$  Hz, 3H), 1.16 (s, 9H), 1.88 (m, 2H), 2.16 (m, 4H), 2.19 (s, 3H), 3.97 (s, 3H), 4.86 (dd,  $J = 5.2$ , 1.6 Hz, 1H), 6.53 (d,  $J = 8.4$  Hz, 1H), 6.88 (m, 2H), 7.14 (dd,  $J = 7.2$ , 1.6 Hz, 1H), 7.73 (d,  $J = 8.8$  Hz, 1H), 7.82 (s, 1H), 7.85 (d,  $J = 8.8$  Hz, 1H), 8.04 (dd,  $J = 6.4$ , 2.0 Hz, 1H), 8.52 (s, 1H). High Res. ES-MS: 511.2824; calc. for  $C_{32}H_{40}O_4+Na$ :

20 511.2825

C. Diastereomeric isomer pair 1 and diastereomeric isomer pair 2 of 6-{1-Ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylic acid methyl ester



25



-40-


Following the procedure described in Example 2; treat 6-{1-[4-(3,3-dimethyl-1-ethyl-2-oxobutoxy)-3-methylphenyl]-1-ethylpropyl}naphthalene-2-carboxylic acid methyl ester with NaBH<sub>4</sub> in THF at 0 C. Chromatograph the residue (CH<sub>2</sub>Cl<sub>2</sub> to 2% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>) to give diastereomeric isomer pair 1 of the title compound as a  
 5 glassy foam (1.004 g, 77%) and diastereomeric isomer pair 2 of the title compound as a glassy foam (102 mg, 8%).

Diastereomeric isomer pair 1: <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 0.64 (t, J = 7.6 Hz, 6H), 0.93 (t, J = 8.0 Hz, 3H), 0.93 (s, 9H), 1.70 (m, 1H), 1.81 (m, 1H), 2.12 (s, 3H), 2.17 (m, 4H), 2.61 (bs, 1H), 3.27 (s, 1H), 3.97 (s, 3H), 4.27 (dd, J = 5.6, 3.6 Hz, 1H),  
 10 6.63 (d, J = 8.8 Hz, 1H), 6.90 (s, 1H), 6.93 (d, J = 8.4 Hz, 1H), 7.18 (d, J = 8.8 Hz, 1H), 7.75 (d, J = 8.8 Hz, 1H), 7.83 (s, 1H), 7.86 (d, J = 8.4 Hz, 1H), 8.04 (dd, J = 7.2, 1.2 Hz, 1H), 8.52 (s, 1H). High Res. ES-MS: 513.2914; calc. for C<sub>32</sub>H<sub>42</sub>O<sub>4</sub>+Na: 513.2981  
 Analysis for C<sub>32</sub>H<sub>42</sub>O<sub>4</sub>; Calcd: C, 78.33; H, 8.63; N, 0.00, Found: C, 78.21; H, 8.75; N,  
 15 0.11.

Diastereomeric isomer pair 2:

<sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 0.64 (t, J = 7.2 Hz, 6H), 0.99 (t, J = 4.4 Hz, 3H), 0.99 (s, 9H), 1.81 (m, 2H), 2.12 (s, 3H), 2.16 (m, 4H), 3.59 (d, J = 4.0 Hz, 1H), 3.97 (s, 3H), 4.35 (m, 1H), 6.67 (d, J = 8.4 Hz, 1H), 6.91 (m, 2H), 7.18 (dd, J = 6.8, 2.0 Hz,  
 20 1H), 7.75 (d, J = 8.8 Hz, 1H), 7.83 (s, 1H), 7.86 (d, J = 8.8 Hz, 1H), 8.05 (dd, J = 6.8, 2.2 Hz, 1H), 8.52 (s, 1H). High Res. ES-MS: 513.2973; calc. for C<sub>32</sub>H<sub>42</sub>O<sub>4</sub>+Na: 513.2981

D. Diastereomeric isomer pair 1 of 6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylic acid  
 25




Add KOH (321 mg, 5.724 mmol) to a mixture of diastereomeric isomer pair 1 of 6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-

-41-

propyl}-naphthalene-2-carboxylic acid methyl ester (702 mg, 1.431 mmol), THF (9 ml), MeOH (3 ml), and H<sub>2</sub>O (1 ml). Heat the reaction mixture to 60 °C and stir it for 14 h. Cool and partition the reaction mixture between Et<sub>2</sub>O and 1N HCl. Dry the organic layer over MgSO<sub>4</sub>, filter, and concentrate the filter in vacuo to give the title

- 5 compound as a glassy solid (665 mg, 97%). <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 0.64 (t, J = 7.6 Hz, 6H), 0.93 (t, J = 8.4 Hz, 3H), 0.93 (s, 9H), 1.72 (m, 1H), 1.82 (m, 1H), 2.12 (s, 3H), 2.19 (q, 4H), 3.28 (s, 1H), 4.28 (dd, J = 6.0, 3.2 Hz, 1H), 6.65 (d, J = 8.4 Hz, 1H), 6.90 (d, J = 2.0 Hz, 1H), 6.94 (dd, J = 6.0, 2.4 Hz, 1H), 7.21 (dd, J = 7.2, 1.6 Hz, 1H), 7.78 (d, J = 8.8 Hz, 1H), 7.86 (s, 1H), 7.89 (d, J = 8.8 Hz, 1H), 8.09 (dd, J = 6.8, 1.6 Hz, 1H), 8.62 (s, 1H). High Res. ES-MS: 499.2832; calc. for C<sub>31</sub>H<sub>40</sub>O<sub>4</sub>+Na: 499.2825
- 10 Analysis for C<sub>31</sub>H<sub>40</sub>O<sub>4</sub>; Calcd: C, 78.11; H, 8.46; N, 0.00, Found: C, 78.03; H, 8.59; N, 0.12.

- 15 E. Diastereomeric isomer pair 1 of 6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylaminoacetic acid methyl ester



- 20 Add glycine methyl ester hydrochloride (95 mg, 0.755 mmol), HOBT (102 mg, 0.755 mmol), EDCI (206 mg, 1.08 mmol), and Et<sub>3</sub>N (0.4 ml, 0.726 mmol) to diastereomeric isomer pair 1 of 6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylic acid (343 mg, 0.72 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (5 ml). Stir the reaction for 14 h. Concentrate the reaction
- 25 mixture in vacuo, and chromatograph the residue (CH<sub>2</sub>Cl<sub>2</sub> to 15% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>) to give the title compound as a colorless foam (373 mg, 95%). <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 0.64 (t, J = 7.6 Hz, 6H), 0.93 (t, J = 8.4 Hz, 3H), 0.93 (s, 9H), 1.68 (m, 1H), 1.81 (m, 1H), 2.12 (s, 3H), 2.18 (q, 4H), 2.60 (bs, 1H), 3.27 (s, 1H), 3.83 (s, 3H),


-42-

4.27 (dd,  $J = 5.6, 3.2$  Hz, 1H), 4.31 (d,  $J = 5.2$  Hz, 2H), 6.63 (d,  $J = 8.0$  Hz, 1H), 6.75 (t,  $J = 4.8$  Hz, 1H), 6.89 (d,  $J = 2.4$  Hz, 1H), 6.93 (dd,  $J = 6.4, 2.0$  Hz, 1H), 7.18 (dd,  $J = 6.8, 2.0$  Hz, 1H), 7.75 (d,  $J = 8.4$  Hz, 1H), 7.83 (d,  $J = 2.0$ , 1H), 7.86 (m, 2H), 8.26 (s, 1H). High Res. ES-MS: 570.3116; calc. for  $C_{34}H_{45}NO_5 + Na$ : 570.3196.

- 5 Analysis for  $C_{34}H_{45}NO_5$ ; Calcd: C, 74.56; H, 8.28; N, 2.56, Found: C, 74.28; H, 8.23; N, 2.68.

F. Diastereomeric isomer pair 1 of 2-[6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylamino]acetic acid


10 acid




- Add 2.5 M LiOH/H<sub>2</sub>O (0.82 ml, 2.04 mmol) to diastereomeric isomer pair 1 of 2-[6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylamino]acetic acid methyl ester (373 mg, 0.681 mmol)
- 15 in THF (3 ml), and MeOH (1.5 ml). Stir the reaction for 14 h; then partition the reaction mixture between Et<sub>2</sub>O and 1N HCl. Dry the organic layer with MgSO<sub>4</sub>, concentrate and triturate the residue with CH<sub>2</sub>Cl<sub>2</sub>/MeOH/hexanes to give the title compound as a white solid (310 mg, 85%). <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>)  $\delta$  ppm: 0.64 (t,  $J = 7.2$  Hz, 6H), 0.93 (t,  $J = 8.0$  Hz, 3H), 0.93 (s, 9H), 1.68 (m, 1H), 1.81 (m, 1H),
- 20 2.11 (s, 3H), 2.15 (q, 4H), 3.28 (s, 1H), 4.27 (dd,  $J = 5.2, 3.6$  Hz, 1H), 4.35 (d,  $J = 4.8$  Hz, 2H), 6.63 (d,  $J = 8.0$  Hz, 1H), 6.83 (t,  $J = 4.8$  Hz, 1H), 6.89 (s, 1H), 6.93 (dd,  $J = 6.4, 2.2$  Hz, 1H), 7.18 (dd,  $J = 7.2, 1.6$  Hz, 1H), 7.74 (d,  $J = 8.8$  Hz, 1H), 7.88 (m, 3H), 8.27 (s, 1H). High Res. ES-MS: 556.3046; calc. for  $C_{33}H_{43}NO_5 + Na$ : 556.3040.

- 25 Example 8. Preparation of diastereomeric isomer pair 1 of N-methyl-2-[6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxyl-amino]acetic acid

-43-



A. Diastereomeric isomer pair 1 of N-methyl-2-[6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carbonyl-amino]acetic acid ethyl ester




5

Following a procedure analogous to Example 7E esterify the diastereomeric isomer pair 1 of 6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carboxylic acid using N-methyl glycine ethyl ester hydrochloride to provide the title compound as a white foam (315 mg, 88%). <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 0.64 (t, J = 7.2 Hz, 6H), 0.93 (t, J = 8.4 Hz, 3H), 0.93 (s, 9H), 1.31 (m, 3H), 1.68 (m, 1H), 1.80 (m, 1H), 2.12 (s, 3H), 2.17 (q, 4H), 2.60 (bs, 1H), 3.11 (s, 2H), 3.16 (s, 1H), 3.27 (s, 1H), 4.05 (s, 0.73H), 4.27 (m, 3H), 4.32 (s, 1.27H), 6.63 (d, J = 8.8 Hz, 1H), 6.90 (s, 1H), 6.93 (d, J = 8.8 Hz, 1H), 7.16 (d, J = 10.0 Hz, 1H), 7.49 (m, 1H), 7.55 (d, J = 7.6 Hz, 1H), 7.68 (m, 1H), 7.81 (s, 1H), 7.87 (m, 2H). High Res. ES-MS: 598.3488; calc. for C<sub>36</sub>H<sub>49</sub>NO<sub>5</sub>+Na: 598.3509. Analysis for C<sub>36</sub>H<sub>49</sub>NO<sub>5</sub>; Calcd: C, 75.10; H, 8.58; N, 2.43, Found: C, 75.04; H, 8.58; N, 2.43.

B. Diastereomeric isomer pair 1 of N-methyl-2-[6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-naphthalene-2-carbonyl-amino]acetic acid ethyl ester

20




-44-

Following a procedure analogous to Example 7F, esterify the diastereomeric isomer pair 1 of N-methyl-2-[6-{1-ethyl-1-[4-(1-ethyl-2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl]-naphthalene-2-carbonyl-amino]acetic acid ethyl ester in THF and EtOH to give the title compound as a white solid (300 mg, quant.).

5 <sup>1</sup>NMR (400MHz, CDCl<sub>3</sub>) δ ppm: 0.64 (t, J = 7.2 Hz, 6H), 0.93 (t, J = 7.6 Hz, 3H), 0.93 (s, 9H), 1.65 (m, 1H), 1.83 (m, 1H), 2.12 (s, 3H), 2.15 (q, 4H), 3.16 (s, 3H), 3.28 (s, 1H), 4.12 (bs, 0.5H), 4.26 (dd, J = 5.2, 3.6 Hz, 1H), 4.36 (s, 1.25H), 6.63 (d, J = 8.4 Hz, 1H), 6.90 (s, 1H), 6.93 (d, J = 8.8 Hz, 1H), 7.17 (d, J = 8.0 Hz, 1H), 7.53 (m, 1H), 7.68 (m, 1H), 7.82 (s, 1H), 7.87 (m, 1H), 7.93 (m, 1H). High Res. ES-MS:  
10 570.3140; calc. for C<sub>34</sub>H<sub>45</sub>NO<sub>5</sub>+Na: 570.3196.


Example 9. 3'-[4-(2-hydroxy-3,3-dimethylbutoxy)-3-methylphenyl]-3'-[12-(t-butoxycarbonylamino)-naphthalene]pentane



15 Treat 3'-[4-(2-Hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-3' [(12-carboxylic acid) -naphthalene] pentane (1 eq.) in CH<sub>2</sub>Cl<sub>2</sub> with, Et<sub>3</sub>N (1.1 eq.) (PhO)<sub>2</sub>PO(N<sub>3</sub>) (1.1 eq) and allow the reaction to stir for about 1 h. Then concentrate the reaction mixture, and add the concentrated solution to a 90°C solution of t-BuOH  
20 and heat with an open stream of nitrogen for about 1.75 h. Cool the reaction to RT, dissolve in a minimal of 1:1 CH<sub>2</sub>Cl<sub>2</sub>:10% EtOAc/Hex, and chromatographed (10% EtOAc/Hex) to give the title compound.

25 Example 10. 3'-[4-(2-Hydroxy-3,3-dimethylbutoxy)-3-methylphenyl]-3'-[12-amino-naphthalene]pentane.

-45-



Treat mixture of 3'-[4-(2-hydroxy-3,3-dimethylbutoxy)-3-methylphenyl]-3'-[12-(t-butoxycarbonylamino)-naphthalene]pentane (1 eq.), anisole (about 20 eq.) in CH<sub>2</sub>Cl<sub>2</sub> (5 ml) with TFA (excess). Stir the reaction for about 2 h, concentrate, and partition the reaction mixture between EtOAc/satd Na<sub>2</sub>CO<sub>3</sub>. Wash the organic layer with water, dry over Na<sub>2</sub>SO<sub>4</sub>, and filter. Concentrate the filtrate and chromatograph the residue (50% CHCl<sub>3</sub>/Hex to CHCl<sub>3</sub>) to give the title compound.

#### Compounds of the Invention – Salts, Stereoisomers, & Prodrugs:

Salts of the compounds represented by Formula I are an additional aspect of the invention. The skilled artisan will also appreciate that the family of compounds of Formula I include acidic and basic members and that the present invention includes pharmaceutically acceptable salts thereof.

In those instances where the compounds of the invention possess acidic or basic functional groups various salts may be formed which are more water soluble and physiologically suitable than the parent compound. Representative pharmaceutically acceptable salts, include but are not limited to, the alkali and alkaline earth salts such as lithium, sodium, potassium, ammonium, calcium, magnesium, aluminum, zinc, and the like. Sodium and potassium salts are particularly preferred. Salts are conveniently prepared from the free acid by treating the acid in solution with a base or by exposing the acid to an ion exchange resin. For example, a carboxylic acid substituent on the compound of Formula I may be selected as -CO<sub>2</sub>H and salts may be formed by reaction with appropriate bases (e.g., NaOH, KOH) to yield the corresponding sodium and potassium salt.

Included within the definition of pharmaceutically acceptable salts are the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention, for example, ammonium, quaternary ammonium, and amine cations,

-46-

derived from nitrogenous bases of sufficient basicity to form salts with the compounds of this invention (see, for example, S. M. Berge, *et al.*, "Pharmaceutical Salts," *J. Phar. Sci.*, 66: 1-19 (1977)). Moreover, the basic group(s) of the compound of the invention may be reacted with suitable organic or inorganic acids to form salts such as acetate,

- 5 benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, camsylate, carbonate, chloride, choline, clavulanate, citrate, chloride, chlorprocaine, choline, diethanolamine, dihydrochloride, diphosphate, edetate, edisylate, estolate, esylate, ethylenediamine, fluoride, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, bromide, chloride, hydrobromide, hydrochloride,
- 10 hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, malseate, mandelate, meglumine, mesylate, mesviate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, palmitate, pamoate, pantothenate, phosphate, polygalacturonate, procane, salicylate, stearate, subacetate, succinate, sulfate, tannate, tartrate, teoclate, tosylate, trifluoroacetate, trifluoromethane
- 15 sulfonate, and valerate.

- Certain compounds of the invention may possess one or more chiral centers and may thus exist in optically active forms. Likewise, when the compounds contain an alkenyl or alkenylene group there exists the possibility of cis- and trans- isomeric forms of the compounds. The R- and S- isomers and mixtures thereof, including racemic
- 20 mixtures as well as mixtures of cis- and trans- isomers, are contemplated by this invention. Additional asymmetric carbon atoms can be present in a substituent group such as an alkyl group. All such isomers as well as the mixtures thereof are intended to be included in the invention. If a particular stereoisomer is desired, it can be prepared by methods well known in the art by using stereospecific reactions with starting materials
- 25 which contain the asymmetric centers and are already resolved or, alternatively by methods which lead to mixtures of the stereoisomers and subsequent resolution by known methods. For example, a chiral column may be used such as those sold by Daicel Chemical Industries identified by the trademarks:
- CHIRALPAK AD, CHIRALPAK AS, CHIRALPAK OD, CHIRALPAK OJ,
- 30 CHIRALPAK OA, CHIRALPAK OB, CHIRALPAK OC, CHIRALPAK OF, CHIRALPAK OG, CHIRALPAK OK, and CHIRALPAK CA-1.

-47-

By another conventional method, a racemic mixture may be reacted with a single enantiomer of some other compound. This changes the racemic form into a mixture of diastereomers. These diastereomers, because they have different melting points, different boiling points, and different solubilities can be separated by conventional means, such as crystallization.


Prodrugs are derivatives of the compounds of the invention which have chemically or metabolically cleavable groups and become by solvolysis or under physiological conditions the compounds of the invention which are pharmaceutically active in vivo. Derivatives of the compounds of this invention have activity in both their acid and base derivative forms, but the acid derivative form often offers advantages of solubility, tissue compatibility, or delayed release in a mammalian organism (see, Bundgard, H., Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985). Prodrugs include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acidic compound with a suitable alcohol, or amides prepared by reaction of the parent acid compound with a suitable amine. Simple aliphatic or aromatic esters derived from acidic groups pendent on the compounds of this invention are preferred prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy) alkyl esters or ((alkoxycarbonyl)oxy)alkyl esters. Particularly preferred esters to use as prodrugs are; methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, morpholinoethyl, and N,N-diethylglycolamido.

N,N-diethylglycolamido ester prodrugs may be prepared by reaction of the sodium salt of a compound of Formula I (in a medium such as dimethylformamide) with 2-chloro-N,N-diethylacetamide (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA; Item No.25,099-6).


Morpholinylethyl ester prodrugs may be prepared by reaction of the sodium salt of a compound of Formula I (in a medium such as dimethylformamide) 4-(2-chloroethyl)morpholine hydrochloride (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA, Item No. C5,220-3). For example, prodrugs may be prepared by reaction of the sodium salt for a compound of Formula I with;



-48-



and sodium iodide to provide the ester prodrug pendent group



Also, lower alkyl (viz., C<sub>1</sub>-C<sub>8</sub>) ester prodrugs may be prepared by conventional means such as reacting the sodium or potassium salt (derived by forming the salt of any acidic compound of the invention; viz., reaction of a base such as KOH with an acidic group such as -CO<sub>2</sub>H) of a compound of Formulae I with an alkyl iodide such as methyl iodide, ethyl iodide, n-propyl iodide, isopropyl iodide.

#### Pharmaceutical Formulations containing the Novel Compounds of the Invention:

Pharmaceutical formulations of the invention are prepared by combining (e.g., mixing) a therapeutically effective amount of the compound of the invention (compounds of Formula I) together with a pharmaceutically acceptable carrier or diluent. The present pharmaceutical formulations are prepared by known procedures using well-known and readily available ingredients.

In making the compositions of the present invention, the compounds of the invention will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), or ointment, containing, for example, up to 10% by weight of the compound. The compounds of the present invention are preferably formulated prior to administration.

-49-

The compounds of the invention may also be delivered by suitable formulations contained in a transderm patch. Alternatively, the compounds of the invention may be delivered to a patient by sublingual administration.

For the pharmaceutical formulations any suitable carrier known in the art can be used. In such a formulation, the carrier may be a solid, liquid, or mixture of a solid and a liquid. Solid form formulations include powders, tablets and capsules. A solid carrier can be one or more substances which may also act as flavoring agents, lubricants, solubilisers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

Tablets for oral administration may contain suitable excipients such as calcium carbonate, sodium carbonate, lactose, calcium phosphate, together with disintegrating agents, such as maize, starch, or alginic acid, and/or binding agents, for example, gelatin or acacia, and lubricating agents such as magnesium stearate, stearic acid, or talc.

In powders the carrier is a finely divided solid which is in admixture with the finely divided Active Ingredient. In tablets a compound of the invention is mixed with a carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about 1 to about 99 weight percent of the compound of this invention. Suitable solid carriers are magnesium carbonate, magnesium stearate, talc, sugar lactose, pectin, dextrin, starch, gelatin, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, low melting waxes, and cocoa butter.

Sterile liquid form formulations include suspensions, emulsions, syrups and elixirs.

The Active Ingredient may be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent or a mixture of both. The compounds can often be dissolved in a suitable organic solvent, for instance aqueous propylene glycol. Other compositions can be made by dispersing the finely divided compounds of the invention in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.

#### Methods of Using the Compounds of the Invention:

-50-

Many disease states are benefited by treatment with the compounds of Formula I include, but are not limited to: disease states characterized by: abnormal calcium regulation, abnormal cell proliferation, abnormal cell differentiation, abnormal immune response, abnormal dermatological conditions, neurodegenerative condition, inflammation, vitamin D sensitivity, and/or hyperproliferative disorders.

Specific disease states benefited by treatment with one or more of the compounds of Formula I include, but are not limited to: Acne, Actinic keratosis, Alopecia, Alzheimer's disease, Benign prostatic hyperplasia, Bladder cancer, Bone maintenance in zero gravity, Bone fracture healing, Breast cancer, Chemoprevention of Cancer, Crohn's disease, Colon cancer, Type I diabetes, Host-graft rejection, Hypercalcemia, Type II diabetes, Leukemia, Multiple sclerosis, Myelodysplastic syndrome, Insufficient sebum secretion, Osteomalacia, Osteoporosis, Insufficient dermal firmness, Periodontal disease, Insufficient dermal hydration, Psoriatic arthritis, Prostate cancer, Psoriasis, Renal osteodystrophy, Rheumatoid arthritis, Scleroderma, Skin cancer, Systemic lupus erythematosus, Skin cell damage from mustard vesicants, Ulcerative colitis, Vitiligo, and Wrinkles.

Particularly preferred is the treatment of psoriasis and/or osteoporosis by administration to a mammal (including a human) of a therapeutically effective amount of compounds of Formula I. By "pharmaceutically effective amount" it is meant that quantity of pharmaceutical agent corresponding to formulae I which prevents, removes or reduces the deleterious effects of a disease state in mammals, including humans.

The specific dose of a compound administered according to this invention to obtain therapeutic or prophylactic effects will, of course, be determined by the particular circumstances surrounding the case, including, for example, the compound administered, the route of administration and the condition being treated. Typical daily doses will contain a pharmaceutically effective amount typically in the range of from about 0.0001 mg/kg/day to about 50 mg/kg/day of body weight of an active compound of this invention. Preferably the dose of compounds of the invention will be from 0.0001 to 5 mg/kg/day of body weight.

Preferably compounds of the invention or pharmaceutical formulations containing these compounds are in unit dosage form for administration to a mammal. The unit dosage form can be a capsule or tablet itself, or the appropriate number of any

-51-

of these. The quantity of Active Ingredient in a unit dose of composition may be varied or adjusted from about 0.0001 to about 1000 milligrams or more according to the particular treatment involved. It may be appreciated that it is necessary to make routine variations to the dosage depending on the age and condition of the patient.

- 5 Dosage will also depend on the route of administration. The compounds of the invention may be administered by a variety of routes including oral, aerosol, rectal, transdermal, sublingual, subcutaneous, intravenous, intramuscular, and intranasal. Particularly preferred is the treatment of psoriasis with an ointment type formulation containing the compounds of the invention. The ointment formulation may be applied  
10 as needed, typically from one to 6 times daily.

Treatment of psoriasis is preferably done with topical application by a formulation in the form of a cream, oil, emulsion, paste or ointment containing a therapeutically effective amount of a compound of the invention. The formulation for topical treatment contains from 0.5 to 0.00005 weight percent, preferably from .05 to  
15 0.0005 weight percent, and most preferably from 0.025 to 0.001 of a Active Ingredient.

For example, two semisolid topical preparations useful as vehicles for VDR modulators in treatment and prevention of psoriasis are as follows:

Polyethylene Glycol Ointment USP (p. 2495)

Prepare Polyethylene Glycol Ointment as follows:

20	Polyethylene Glycol 3350	400 g.
	Polyethylene Glycol 400	<u>600 g.</u>
	To make	1000 g.

Heat the two ingredients on a water bath to 65°C. Allow to cool, and stir until congealed. If a firmer preparation is desired, replace up to 100 g of the polyethylene  
25 glycol 400 with an equal amount of polyethylene glycol 3350.

Hydrophilic Ointment USP (p. 1216)

Prepare Hydrophilic Ointment as follows:

	Methylparaben	0.25 g.
	Propylparaben	0.15 g.
30	Sodium Lauryl Sulfate	10 g.
	Propylene Glycol	120 g.
	Stearyl Alcohol	250 g.

-52-

White Petrolatum	250 g.
Purified Water	<u>370 g.</u>
To make about	1000 g.

5 The Stearyl Alcohol and White Petrolatum are melted on a steam bath, and warmed to about 75°C. The other ingredients, previously dissolved in the water are added, warmed to 75°C, and the mixture stirred until it congeals.

For each of the above formulations the Active Ingredient is added during the heating step in an amount that is from 0.5 to 0.00005 weight percent, preferably from .05 to 0.0005 weight percent, and "USP" most preferably from 0.025 to 0.001 weight percent  
10 of the total ointment weight. (Source: - United States Pharmacopoeia 24, United States Pharmacopeial Convention, 1999)

Conventional therapy for osteoporosis includes; (i) estrogens, (ii) androgens, (iii) calcium supplements, (iv) vitamin D metabolites, (v) thiazide diuretics, (vi) calcitonin, (vii) bisphosphonates, (viii) SERMS, (ix) fluorides and (x) Parathyroid hormone (PTH)  
15 (see, Harrison's Principles of Internal Medicine, 13<sup>th</sup> edition, 1994, published by McGraw Hill Publ., ISBN 0-07-032370-4, pgs. 2172-77; the disclosure of which is incorporated herein by reference). Any one or a combination of these conventional therapies may be used in combination with the method of treatment using compounds of Formula I as taught herein. For example, in a method of treating osteoporosis, the vitamin D receptor  
20 modulator compounds of the invention may be administered separately or simultaneously with a conventional therapy. Alternatively, the vitamin D receptor modulator compounds of the invention may be combined with conventional therapeutic agents in a formulation for treatment of osteoporosis such as set out below:

A formulation for treating osteoporosis comprising:

25 Ingredient (A1): a vitamin D receptor modulator represented by Formula (I), or a pharmaceutically acceptable salt or prodrug derivative thereof;

Ingredient (B1): one or more co-agents that are conventional for treatment of osteoporosis selected from the group consisting of: estrogens, androgens, calcium supplements, vitamin D metabolites, thiazide  
30 diuretics, calcitonin, bisphosphonates, SERMS, fluorides, and PTH

Ingredient (C1): optionally, a carrier or diluent.

-53-

Typically useful formulations are those wherein the weight ratio of (A1) to (B1) is from 10:1 to 1:1000 and preferably from 1:1 to 1:100.

Combination Therapy for Psoriasis:

5           Conventional therapy for psoriasis includes topical glucocorticoids, salicylic acid, crude coal tar, ultraviolet light, and methotrexate (see, Harrison's Principles of Internal Medicine, 13<sup>th</sup> edition, 1994, published by McGraw Hill Publ., ISBN 0-07-032370-4, pgs. 2172-77). Any one or combination of these conventional therapies may be used in combination with the method of treatment using compounds of Formula I as taught  
10       herein. For example, in a method of treating psoriasis, the vitamin D receptor modulator compounds of the invention (e.g., as defined by Formula I) may be topically administered separately or simultaneously with a conventional therapy. Alternatively, the vitamin D receptor modulator compounds of the invention may be combined with conventional therapeutic agents in a topically applied formulation for treatment of psoriasis such as set  
15       out below:

A formulation for treating psoriasis comprising:

Ingredient (A2): a vitamin D receptor modulator represented by Formula(I), or a pharmaceutically acceptable salt or prodrug derivative thereof;

20           Ingredient (B2): one or more co-agents that are conventional for treatment psoriasis selected from the group consisting of: topical glucocorticoids, salicylic acid, or crude coal tar.

Ingredient (C2): optionally, a carrier or diluent.

Typically useful formulations are those wherein the weight ratio of (A2) to (B2) is from  
25       1:10 to 1:100000 and preferably from 1:100 to 1:10000.

-54-

Experimental Results:

Table 2  
Summary of Experimental Results

Test Cmpd. <sup>1</sup>	RXR-VDR heterodimer <sup>2</sup> EC <sub>50</sub> (nM)	VDR EC <sub>50</sub> (nM) (Caco-2 cells) <sup>3</sup>	OCN Promoter <sup>4</sup> EC <sub>50</sub> (nM)	Mouse Hypercal <sup>5</sup> μg/Kg/d
Ex. 1	40	194	2.6	1000
Ex. 2A	30.2	466	6.0	300
Ex. 2B	25.3		18.4	1000
Ex. 3	0.4	35	0.1	<300
Ex. 4	105	550	41	1000
Ex. 5	1	162	0.2	<30
Ex. 6	715	714	104	
Ex. 7	2.6		0.7	100
Ex. 8	2.0		0.6	300

5

Table 3  
Summary of Experimental Results

Test Cmpd. <sup>1</sup>	Keranocyte Proliferation IC <sub>50</sub> (nM)	IL-10. IC <sub>50</sub> (nM)
Ex. 1	653	76

-55-

Ex. 2	47	49
Ex. 3	2	
Ex. 4	>1000	
Ex. 5	24	5

Explanation of Tables 2 and 3 column numerical superscripts:

1. Test Compound numbers refer to the products of the corresponding Example Nos. that is, compounds within the scope of the invention.

5        2. The RXR-VDR heterodimerization (SaOS-2 cells) test is described in the "Assay" section of the Description, *infra*.

3. The VDR CTF (Caco-2 cells) test is described in the "Assay" section of the Description, *infra*.

10       4. The OCN Promoter test is described in the "Assay" section of the Description, *infra*.

5. The Mouse Hypercalcemia test is described in the "Assay" section of the Description, *infra*.

6. The keratinocyte proliferation assay is described in the "Assay" section of the Description, *infra*.

15       7. The IL-10 induction assay is described in the "Assay" section of the Description, *infra*.

### Assay Methods

#### Use of the Assay Methods:

20       The evaluation of the novel compounds of the invention for osteoporosis and other related diseases is done using a plurality of test results. The use of multiple assays is beneficial since it is preferred that the combined properties of (i) high activity for the vitamin D receptor, and (ii) prevention of hypercalcemia be achieved to effect treating diseases, which are also aspects of this invention. Some of the tests described below are  
25       believed related to other tests and measure related properties of compounds.

Consequently, a compound may be considered to have utility in the practice of the invention if it meets at least one preferably two or more, if not all, of the acceptance criteria for the above described tests.



-56-

The evaluation of the novel compounds of the invention for psoriasis is done using the Keratinocyte Proliferation Assay in combination with other assays that measure inhibition of IL-2 production and stimulation of IL-10 production in peripheral blood mononuclear cells (PBMCs).

5 Brief Description, Utility and Acceptance Criteria for the Assay Methods:

1. The RXR-VDR heterodimer Assay:

This assay provides the VDR activity of a test compound. It is desirable to have low EC<sub>50</sub> values for a compound in this assay. The lower the EC<sub>50</sub> value, the more  
10 active the compound will be as a VDR agonist. Desired assay results are EC<sub>50</sub> values less than or equal to 600 nM. Preferred assay results are less than 250 nM, and most preferably less than 150 nM.

(1) Materials and Method for RXR-VDR Heterodimerization Assay Transfection

- 15 Method: Reagents: FuGENE 6 Transfection Reagent (Roche Cat # 1 814 443);  
Growth Media: D-MEM High Glucose (Gibco BRL Cat # 11054-020), 10% heat inactivated FBS, (Gibco BRL Cat# 10092-147) 1% antibiotic-antimycotic (Ab-Am); (Gibco BRL Cat # 15240-062 ).
- 20 Cells: Grow SaOS-2 cells in T-150 cm<sup>2</sup> culture flasks in *growth media* keeping the density at 5-6 x 10<sup>5</sup> cells/ml. Passage cells 1:3 twice a week. Add Trypsin EDTA (Gibco BRL Cat # 25300-020) and incubate. Resuspend cells in plating media and transfer into growth media.
- 25 Wash Media: HBSS Low Glucose Without Phenol Red (Gibco BRL Cat # 14175-095), 1% Ab-Am. Plating Media: D-MEM Low Glucose Without Phenol Red (Gibco BRL Cat # 11054-020), 1% Ab-Am; D-MEM; 10 % Stripped FBS (Hyclone Cat# SH30068.03 Lot # AHM9371);
- 30 Transfection / Treatment Media: D-MEM Low Glucose Without Phenol Red only; T-150 cm<sup>2</sup> culture flask: Use Corning Coastar T-150 cm<sup>2</sup> culture flask (Cat # 430825) to grow the cells.

-57-

Luciferase Assay Reagent: Use Steady-Glo Luciferase Reagent from Promega (Cat # E2550) Consists of: E2533 Assay Substrate, lyophilized product and E2543 Assay Buffer. Thaw at room temperature and store.

- 5 Cell Harvesting/Count: Aspirate media from culture flask, rinse cells with HBSS and aspirate. Add trypsin and incubate. When cells appear detached, resuspend cells in *growth media*. Transfer into a new flask with fresh *growth media* for passaging the cells. Plate 96 well plates and two extra plates. Mix the cell suspension using pipette. To count the cells using a *Hematocytometer*.

10

Plate seeding: Use plating media 10 % Stripped FBS in D-MEM Low Glucose, without Phenol Red, 1% Ab-Am. Plate 14 plates @ 165  $\mu$ l / well. In sterile flask add cell suspension to plating media and mix. Add cells / well. Place the cells in the incubator.

- Cells should be about 75 % confluent prior to transfection. DAY 2: Transfection: Step 1, DNA and Media: Add plain DMEM media to tubes for mixing the DNA; add the Reporter gene pFR-LUC; and add the Gal4-RXR-DEF and VP16-VDR-LBD. Step 2, FuGENE and Media: Prepare plain DMEM media in a tubes for mixing FuGENE, add *FuGENE 6 Transfection Reagent*, and incubate. Step 3, FuGENE, DNA and Media Complex: Add FuGENE Media complex from step 2 to DNA Media complex from step 1  
15 and incubate. Step 4, FuGENE, DNA and Media Complex to 96 well plate: Add FuGENE-DNA-Media complex from step 3 to each plate. Incubate.

20

Day 3: Dosing: Treatment preparation . Allow for transfection time.

Make a stock solution of the compounds in DMSO and vortex until all the compounds  
25 have been dissolved. Further dilute in D-MEM (Low Glucose – without Phenol Red)  
Add compounds in quadruplicate to give desired final volume then incubate.

- Day 4: Luciferase Assay: Read the plates after drug treatment. Remove part of media from all the wells and leave remainder. Add Steady-Glo Luciferase Reagent mixture/  
30 wells and incubate. Count each well using a Luminescence counter, Top Count NXT by Packard preferably set a delay between plates to reduce the background.

-58-

The Caco-2 cell Co-transfection Assay:

The Caco-2 cell assay is an indicator for the undesirable condition of hypercalcemia. This co-transfection assay is a surrogate assay for in vivo calcemic activity of VDR ligands. It is desirable to have high EC<sub>50</sub> values for a test compound in this assay. The higher the EC<sub>50</sub> values for a compound the less calcemic it will be in vivo. Desired assay results are EC<sub>50</sub> greater than or equal to 300 nM. Preferred assay results are greater than 1000 nM.

Caco-2 cells, grown in phenol red free, DMEM (Invitrogen, Carlsbad, CA) containing 10 % charcoal-stripped FBS (Hyclone, Logan, UT), are transfected with Fugene 6 reagent (Roche Diagnostics, Indianapolis, IN). Cells (5000/well) are plated 18 h before transfection in a 96 well plate. The cells are transfected with Gal4-responsive reporter pFRLuc (150 ng, Stratagene, La Jolla CA) and the receptor expression vector pGal4-VDR-LBD (10 ng), along with Fugene 6 reagent (0.2 µl/well). The DNA-Fugene complex is formed by incubating the mixture for 30 m at room temperature. The cells are transfected in triplicate for 5 h, and treated with various concentrations of VDR ligands (from 0.01 nM to 10,000 nM concentration range) 18h post-transfection. The luciferase activity is quantified using Steady-Glo reagent kit (Promega, Madison, WI) as per manufacturer's specifications.

The OCN (osteocalcin) Promoter Assay

The OCN Promoter Assay is an indicator and marker for osteoporosis. Desired assay results are EC<sub>50</sub> less than or equal to 325 nM. Preferred assay results are less than 50 nM.

The activation of osteocalcin by VDR ligands is evaluated in a rat osteoblast-like cell line RG-15 (ROS 17/2.8) stably expressing rat osteocalcin promoter fused with luciferase reporter gene. The stable cell lines are established as reported before (Activation of Osteocalcin Transcription involves interaction of protein kinase A- and Protein kinase C-dependent pathways. Boguslawski, G., Hale, L. V., Yu, X.-P., Miles, R. R., Onyia, J. E., Santerre R. F., Chandrasekhar, S. J Biol. Chem. 275, 999-1006, 2000).

Confluent RG-15 cells maintained in DMEM/F-12 medium (3:1) containing 5% FBS, 300 µg/ml G418 and at 37°C under 5% CO<sub>2</sub>/95% air atmosphere are trypsinized (0.25% trypsin) and plated into white opaque 96-well cell culture plates (25000 cells/well). After

-59-

24 hr, cells (in DMEM/F-12 medium + 2% FBS) are treated with various concentrations of compounds, dissolved in DMSO. The final DMSO concentration remains at 0.01% (v/v). After 48 hr treatment, the medium is removed, cells are lysed with 50 µl of lysis buffer (From Luciferase reporter assay system, Roche Diagnostics, Indianapolis, IN) and then assayed for luciferase activity using the Luciferase Reporter Gene Assay kit from Boehringer Mannheim as per manufacturer's specifications.

#### The Mouse Hypercalcemia Assay

The Mouse Hypercalcemia Assay is a six day hypercalcemia test for toxicity and selectivity. Acceptable test results are levels greater than 30 µg/kg/day. Preferred assay results are levels greater than 300 µg/kg/day.

Weanling, virus -antibody-free, five to six weeks old female DBF mice (Harlan, Indianapolis, IN) are used for all the studies. Animals are allowed to acclimate to local vivarium conditions for 2 days. Mice are maintained on a 12 hr light/dark cycle at 22°C with ad lib access to food (TD 5001 with 1.2% Ca and 0.9% P, Teklad, Madison, WI) and water. The animals then are divided into groups with 4-5 mice per group. Different doses of test compounds prepared in 10% ethanol and 90% sesame oil, or in an aqueous suspension of sodium lauryl sulfate and CMC (the latter formulation for acidic compounds) are administered to mice orally via gavage for 6 days. 1α-25(OH)<sub>2</sub>D<sub>3</sub> 0.5µg/kg/d was also given to one group of mice as the positive control. Serum ionized calcium is evaluated at 6 hours after the last dosing under isoflurane anesthesia by Ciba-Corning Ca++/PH Analyzer, (Model 634, Chiron Diagnostics Corp., East Walpole, MA). Raw data of group differences is assessed by analysis of variance (ANOVA) using Fisher's protected least significant difference (PLSD) where the significance level was P< 0.05. The highest dose that did not cause hypercalcemia, as defined by the 97.5% reference distribution of the control population, is considered "the no effect level".

#### The Keratinocyte Proliferation Assay

This Assay is indicative for the treatment of psoriasis. An acceptable test result is IC<sub>50</sub> value of less than or equal to 300 nM. Preferred assay results are IC<sub>50</sub> values of less than 100 nM.

-60-

KERtr cells (Human skin keratinocyte are transformed with a retrovirus vector, obtained from ATCC, then are plated in 96-well flat-bottomed plates (3000 cells/well) in 100  $\mu$ l keratinocyte serum free medium supplemented with bovine pituitary extract in the absence of EGF (Life Technologies, Rockville, MD) and are incubated at 37°C for two

5 days. The cells are treated with various concentrations of VDR ligands (ten-fold serial dilution from 10,000 nM to 0.1 nM in triplicate), dissolved in 100  $\mu$ l keratinocyte serum free medium supplemented with bovine pituitary extract in the absence of EGF and are incubated at 37°C for 72 hr. BrdU (5-bromo-2'-deoxyuridine) incorporation is analyzed as a measure of DNA replication (Cell proliferation ELISA kit, Roche Diagnostics,

10 Indianapolis, IN) and absorbance is measured at 405 nm. Potency values ( $IC_{50}$ ) values were determined as the concentration (nM) of compound that elicited a half-maximal response.

#### The IL-10 induction Assay

15 This is an in vitro efficacy assay for psoriasis, abscess and adhesion. Psoriasis involves both keratinocytes and immune cells. IL-10 is a unique cytokine because it is anti-inflammatory and immunosuppressive. This assay tells us whether a VDRM is able to function as an agonist in PBMCs (primary blood mononuclear cells) or not. A lower EC50 value is desirable in this assay since a compound with a lower EC50 value will be a

20 better agonist in PBMCs. An acceptable test result is an EC50 value of less than 200 nM. Preferred assay results are EC50 values of less than 100 nM.

#### Isolation of peripheral blood mononuclear cells (PBMCs):

Collect 50 ml of human blood and dilute with media, RPMI-1640. Add diluted

25 blood to sterile tubes with ficol. Centrifuge the tubes. Discard the top layer and collect the cells from middle layer. Divide all cells into four tubes and add media. Centrifuge. Aspirate off media and resuspend the cells. Collect all cells. Centrifuge at 1200 rpm for 10 m. Resuspend the cells in RPMI-1640 with 2% FBS and then count cells.

Stimulation of PBMC: Prepare TPA in DMSO. Dissolve PHA in water. Plate

30 TPA/PHA treated PBMCs in well plates. Incubate the cells.

-61-

Treatment: Prepare all compound dilutions in plain RPMI- 1640 media. Add diluted compound and incubate. Sample Collection and assay: Remove all the cells by centrifugation and assay the supernatant for IL-10 by immunoassay using anti-human IL-10 antibody coated beads, as described by the manufacturer (Linco Research Inc., St. Charles, MO).

#### Other Compound Assay Standards

An alternative measure of the therapeutic index (bone efficacy vs Hypercalcemia) of compounds of the invention for treatment of osteoporosis is a numerical ratio calculated as follows:

**Dose Threshold needed to induce hypercalcemia  
divided by**

**Dose Threshold needed for bone efficacy**

An alternative measure of the therapeutic index (in vivo keratinocyte proliferation vs. hypercalcemia) of compounds of the invention for treatment of psoriasis is a numerical ratio calculated as follows:

**Dose Threshold needed to induce hypercalcemia  
divided by**

**Dose Threshold needed to induce keratinocyte proliferation**

For the above ratios, Dose Thresholds are determined from dose response curve data.

#### The CaT1 (calcium transporter 1) Assay

The CaT1 Assay is an indicator for the undesirable condition of hypercalcemia. The higher the EC50 values for a compound the less calcemic it will be in vivo. Desired assay results are EC50 greater than or equal to 500 nM. Preferred assay results are greater than 1000 nM.

Human colon carcinoma, Caco-2 cells, maintained in DMEM (high glucose with 25 mM Hepes buffer; Invitrogen, Carlsbad, CA) supplemented with 10 % FBS (Invitrogen, Carlsbad, CA), are plated at 5500 cell per well in a 96-well plate in a total volume of 100 µl/well. The cells are kept in the 96-well plate for 6 days to differentiate them to small intestinal cells that express the calcium transporter, CaT1. On day 3 after plating, old media is removed and replaced with fresh media (150 µl/well). On day 6 the

-62-

old media is removed and the cells are kept in treatment media (180 µl/well) that contained 10 % charcoal stripped FBS (Hyclone, Logan, UT) in DMEM (low glucose, without phenol red; Invitrogen, Carlsbad, CA). The cells are treated with various concentrations of VDR ligands (from 0.01 nM to 10,000 nM concentration range)

5 prepared in treatment media (20 µl/well). Twenty hours post-treatment, total RNA is prepared by RNeasy 96 method as described by the manufacturer (Qiagen, Valencia, CA). The RNA is reverse transcribed and amplified for human CaT1 and GAPDH (control) messages by quantitative RT-PCR using ABI PRISM 7900HT Sequence Detection System according to manufacturer's instructions (Applied Biosystems, Foster


10 City, CA). Optimized primer pairs and probes for human CaT1 and GAPDH genes are obtained commercially (Applied Biosystems, Foster City, CA). Each 20 µl quantitative RT-PCR reaction in a 384-well Taqman PCR plate consists of forward and reverse primers (900 nM), Taqman probe (200 nM), total RNA (4 µl from each well of the 96-well culture plate) and 10 µl of Taqman Universal PCR Master Mix (Roche Diagnostics,

15 Indianapolis, IN). Reactions are incubated at 48°C for 30 m, followed by 10 m at 95°C and subjected to 40 cycles of PCR (95°C for 15 seconds followed by 60°C for 1 m). GAPDH is used as an internal control and its primer and probe set are obtained commercially (Applied Biosystems, Foster City, CA).

-63-

## CLAIMS:

1. A compound or a pharmaceutically acceptable salt or a prodrug derivative thereof represented by Formula(I):



wherein

R and R' are independently C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, or together R and R' form a substituted or unsubstituted, saturated or unsaturated cycloalkyl ring having from 3 to 8 carbon atoms;

10

RP<sub>3</sub> and RN are independently selected from the group consisting of hydrogen, halo, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, -O-C<sub>1</sub>-C<sub>5</sub> alkyl, -S-C<sub>1</sub>-C<sub>5</sub> alkyl, -O-C<sub>1</sub>-C<sub>5</sub> haloalkyl, -CN, -NO<sub>2</sub>, acetyl, -S-C<sub>1</sub>-C<sub>5</sub> haloalkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, and C<sub>3</sub>-C<sub>5</sub> cycloalkenyl;

RP is selected from the group consisting of: hydrogen, halo, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, -O-C<sub>1</sub>-C<sub>5</sub> alkyl, -S-C<sub>1</sub>-C<sub>5</sub> alkyl, -O-C<sub>1</sub>-C<sub>5</sub> haloalkyl, -CN, -NO<sub>2</sub>, acetyl, -S-C<sub>1</sub>-C<sub>5</sub> haloalkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, and C<sub>3</sub>-C<sub>5</sub> cycloalkenyl;

15

(L<sub>P1</sub>), (L<sub>P2</sub>), and (L<sub>NP</sub>) are divalent linking groups independently selected from the group consisting of: a bond, -(CH<sub>2</sub>)<sub>m</sub>-C(OH)-, -(CH<sub>2</sub>)<sub>m</sub>-O-, -(CH<sub>2</sub>)<sub>m</sub>-S-, -(CH<sub>2</sub>)<sub>m</sub>-S(O)-, -(CH<sub>2</sub>)<sub>m</sub>-S(O)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>m</sub>-N(R<sub>40</sub>)-, -(CH<sub>2</sub>)<sub>m</sub>-C(R<sub>40</sub>)(R<sub>41</sub>)-, -(CH<sub>2</sub>)<sub>m</sub>-C(O)-, -N(R<sub>40</sub>)-C(O)-, -(CH<sub>2</sub>)<sub>m</sub>-CH=CH-, and -(CH<sub>2</sub>)<sub>m</sub>-C≡C-;

20

where m is 0-5;

R<sub>40</sub> and R<sub>41</sub> each is independently selected from the group consisting of: hydrogen, C<sub>1</sub>-C<sub>5</sub> alkyl, C<sub>1</sub>-C<sub>5</sub> hydroxyalkyl, C<sub>1</sub>-C<sub>5</sub> haloalkyl, C<sub>2</sub>-C<sub>5</sub> alkenyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, C<sub>3</sub>-C<sub>5</sub> cycloalkenyl;

25



-64-

$Z_P$  is selected from the group consisting of: branched  $C_3$ - $C_5$  alkyl,  $C_3$ - $C_{10}$  hydroxyalkyl,  $C_3$ - $C_{10}$  hydroxyalkenyl,  $C_3$ - $C_{10}$  hydroxyalkynyl,  $C_3$ - $C_{10}$  hydroxycycloalkyl,  $C_4$ - $C_{10}$  hydroxy cycloalkenyl, and oxocycloalkyl;

$Z_{NP}$  is selected from the group consisting of:  $C_1$ - $C_5$  alkyl,  $C_2$ - $C_5$  alkenyl,  $C_3$ - $C_5$  cycloalkyl,  $C_3$ - $C_5$  cycloalkenyl,  $C_1$ - $C_5$  hydroxyalkyl,  $C_1$ - $C_5$  haloalkyl,  $C_1$ - $C_5$  alkylaryl,  $C_1$ - $C_5$  hydroxyalkylaryl,  $C_0$ - $C_5$  alkyl- $CO_2H$ ,  $C_0$ - $C_3$  alkyl-cycloalkyl- $CO_2H$ ,  $C_0$ - $C_5$  alkyl- $N(R_{40})(R_{41})$ ,  $-X-(C_1-C_5 \text{ alkyl})$ ,  $-X-(C_1-C_5 \text{ alkenyl})$ ,  $-X-(C_3-C_5 \text{ cycloalkyl})$ ,  $-X-(C_3-C_5 \text{ cycloalkenyl})$ ,  $-X-(C_1-C_5 \text{ haloalkyl})$ ,  $-X-(C_1-C_5 \text{ hydroxyalkyl})$ ,  $-X-(C_1-C_5 \text{ alkylaryl})$ ,  $-X(OC_1-C_5 \text{ alkyl})$ ,  $-XN(R_{40})(R_{41})$ ,  $-XN(R_{40})\text{aryl}$ ,  $-N(CH_3)(OCH_3)$ ,  $-N(OH)(CH_3)$ ,  $-N(R_{42})-(C_1-C_5 \text{ alkyl})CO_2H$ ,  $-N(R_{42})-(C_1-C_5 \text{ alkyl})C(O)(C_1-C_5 \text{ alkyl})$ ,  $-N(R_{42})-(C_1-C_5 \text{ alkyl})C(O)(OC_1-C_5 \text{ alkyl})$ ,  $-N(R_{42})-SO_2-(C_1-C_5 \text{ alkyl})$ ,  $-NR(42)-S(O)-(C_1-C_5 \text{ alkyl})$ ,  $-P(O)-(OC_1-C_5 \text{ alkyl})_2$ , heteroalkyl, heteroaryl, and  $-N=C(R_{40})N(R_{40})(R_{41})$ ;

$R_{42}$  is selected from the group consisting of: H,  $C_1$ - $C_3$  alkyl, and  $C_1$ - $C_3$  haloalkyl; and

$X$  is selected from the group consisting of: O,  $C(O)$ ,  $C(S)$ ,  $S(O)$ , and  $SO_2$ ; provided that  $-(L_{NP})-Z_{NP}$  is substituted at either the 12 or 13 position of the naphthalene ring; or a pharmaceutically acceptable salt, solvate, prodrug, enantiomer, racemate, diastereomer or mixture of diastereomers thereof.

20

2. A compound of Claim 1

wherein

$R$  and  $R'$  are independently methyl or ethyl;

$RP$  is hydrogen or methyl;

$RP_3$  and  $RN$  are independently hydrogen, methyl, ethyl,  $-O$ -methyl, or cyclopropyl;

$(L_{P1})$  is a bond;

$(L_{P2})$  is a bond,  $-CH_2-$ ,  $-CH(OH)-$ , or  $-C(Me)OH-$ ;

$(L_{NP})$  is a bond,  $-C(O)-$ ,  $-C(O)NH-$ , or  $-C(O)N(Me)-$ ;

$Z_P$  is selected from the group consisting of: 1,1-dimethylethyl, 1-

30


-65-

hydroxycyclopentyl, 1-hydroxycyclohexyl, 3-ethyl-3-hydroxypentyl, 3-ethyl-3-hydroxypentenyl, and 3-ethyl-3-hydroxypentynyl;


$Z_{NP}$  is  $-CO_2H$ ,  $-CO_2(R40)$ ,  $-N(R40)(R41)$ ,  $-NH-C(Me)(OH)-C(O)OH$ ,  
 $-C(O)NMe-CH_2-C(O)OH$ ,  $-C(O)NMe-CH_2-C(O)OMe$ ,  $-C(O)NMe-CH_2-C(O)OEt$ ,  
 5  $-C(O)NMe-CH_2-C(O)OiPr$ ,  $-C(O)NMe-CH_2-C(O)tBu$ ,  $-cyclopropyl-C(O)OH$ ,  
 $-cyclobutyl-C(O)OH$ ,  $-C(O)NMe-C(Me)_2-C(O)OH$ ,  $-C(O)N(R40)S(O)(R42)$ ,  
 $-C(O)N(R40)SO_2R42$ ,  $-C(O)-N(R40)-5-tetrazolyl$ ,  $-C(O)N(R40)-(C_1-C_5 \text{ alkyl})-$   
 $S(O)R42$ ,  $-C(O)N(R40)-(C_1-C_5 \text{ alkyl})-S(O)_2R42$ , and  $-CH_2CO_2H$ .

- 10 3. A compound represented by Formula(C1) to (C8) or a pharmaceutically acceptable salt or prodrug derivative thereof:

C1)




C2)




15

C3)




-66-

C4)




C5)




5


C6)



C7)



C8)



10

-67-

wherein R1 is H, methyl, or ethyl; R2 is H or methyl; R3 is H, methyl, or ethyl, and R4 is H or methyl.

- 5           4.       The compound of claim 3 represented by formulae (C1) to (C10) or a pharmaceutically acceptable salt or prodrug derivative thereof: where R1 is a methyl, or ethyl; and R2 is H or methyl.
5.       The compound of claim 3 represented by the structural Formula(C2) or a  
10   pharmaceutically acceptable salt thereof.
6.       A compound represented by the structural Formula(C1) or a pharmaceutically acceptable salt thereof.
- 15           7.       A prodrug derivative of the compound of any of claims 1 to 3 wherein the prodrug is a methyl ester; ethyl ester; N,N-diethylglycolamido ester; or morpholinylethyl ester.
8.       A salt derivative of the compound of any of claims 1 to 5 wherein the salt  
20   is sodium or potassium.
9.       A pharmaceutical formulation comprising the compound of any of claims 1 to 7 either with a pharmaceutically acceptable carrier or diluent.
- 25           10.       A formulation for treating osteoporosis comprising:  
a compound of Formula I of any of claims 1 to 8;  
one or more co-agents selected from the group consisting of: estrogens,  
androgens, calcium supplements, vitamin D metabolites, thiazide diuretics, calcitonin,  
bisphosphonates, SERMS, fluorides; and  
30   optionally, a carrier or diluent.

-68-

11. The formulation of claim 9 wherein the weight ratio of a compound of Formula I and the one or more co-agents is from 10:1 to 1:1000.

12. A formulation for treating psoriasis comprising:  
5 a compound of Formula I of any of claims 1 to 8;  
one or more co-agents selected from the group consisting of: topical glucocorticoids, salicylic acid, crude coal tar; and optionally, a carrier or diluent.

10 13. The formulation of claim 12 wherein the weight ratio of Formula I and the one or more co-agents is from 1:10 to 1:100000.

14. Use of compound of Formula 1 in any of claims 1-8 in the manufacture of a medicament for the treatment of osteoporosis.

15 15. A pharmaceutical composition comprising a compound of Formula 1 in any of claims 1-8 for treating osteoporosis.

16. A method of treating a mammal to prevent or alleviate the pathological  
20 effects of Acne, Actinic keratosis, Alopecia, Alzheimer's disease, Bone maintenance in zero gravity, Bone fracture healing, Breast cancer, Chemoprevention of Cancer, Crohn's disease, Colon cancer, Type I diabetes, Host-graft rejection, Hypercalcemia, Type II diabetes, Leukemia, Multiple sclerosis, Myelodysplastic syndrome, Insufficient sebum secretion, Osteomalacia, Osteoporosis, Insufficient dermal firmness, Insufficient dermal  
25 hydration, Psoriatic arthritis, Prostate cancer, Psoriasis, Renal osteodystrophy, Rheumatoid arthritis, Scleroderma, Skin cancer, Systemic lupus erythematosus, Skin cell damage from, mustard vesicants, Ulcerative colitis, Vitiligo, or Wrinkles; wherein the method comprises administering a pharmaceutically effective amount of at least one compound according to any one of claims 1 to 8.

30 17. The method of claim 16 for the treatment of psoriasis.

-69-

18. The method of claim 16 for the treatment of osteoporosis.

19. A method of claim 16 for treating a mammal to prevent or alleviate skin cell damage from mustard vesicants.

5

20. A method of treating a mammal to prevent or alleviate the pathological effects of benign prostatic hyperplasia or bladder cancer wherein the method comprises administering a pharmaceutically effective amount of at least one compound according to any one of claims 1 to 8.

10

21. A method of treating or preventing disease states mediated by the Vitamin D receptor, wherein a mammal in need thereof is administered a pharmaceutically effective amount of the compound according to any one of Claims 1 to 8.

15

22. A compound as claimed in any one of Claims 1 to 8 for use in treating a mammal to prevent or alleviate the pathological effects of Acne, Actinic keratosis, Alopecia, Alzheimer's disease, Bone maintenance in zero gravity, Bone fracture healing, Breast cancer, Chemoprevention of Cancer, Crohn's disease, Colon cancer, Type I diabetes, Host-graft rejection, Hypercalcemia, Type II diabetes, Leukemia, Multiple sclerosis, Myelodysplastic syndrome, Insufficient sebum secretion, Osteomalacia, Osteoporosis, Insufficient dermal firmness, Insufficient dermal hydration, Psoriatic arthritis, Prostate cancer, Psoriasis, Renal osteodystrophy, Rheumatoid arthritis, Scleroderma, Skin cancer, Systemic lupus erythematosus, Skin cell damage from, mustard vesicants, Ulcerative colitis, Vitiligo, or Wrinkles.

25

23. A compound as claimed in any one of Claims 1 to 8 for use in treating or preventing disease states mediated by the Vitamin D receptor.

24. A compound as claimed in any one of Claims 1 to 8 for use in treating a mammal to prevent or alleviate the pathological effects of benign prostatic hyperplasia or bladder cancer.

30

-70-

25. A compound as claimed in any one of Claims 1 to 8 substantially as hereinbefore described with reference to any of the Examples.

26. A process for preparing a compound as claimed in any one of claims 1 to  
5 8 substantially as hereinbefore described with reference to any of the Examples.

27. The use of a compound as claimed in any one of claims 1 to 8 substantially as herein described with reference to any of the Assays and Tables for mediating the Vitamin D receptor.